

## THE PRICE OF STABILITY OF WEIGHTED CONGESTION GAMES\*

GEORGE CHRISTODOULOU<sup>†</sup>, MARTIN GAIRING<sup>†</sup>, YIANNIS GIANNAKOPOULOS<sup>‡</sup>,  
AND PAUL G. SPIRAKIS<sup>§</sup>

**Abstract.** We give exponential lower bounds on the Price of Stability (PoS) of weighted congestion games with polynomial cost functions. In particular, for any positive integer  $d$  we construct rather simple games with cost functions of degree at most  $d$  which have a PoS of at least  $\Omega(\Phi_d)^{d+1}$ , where  $\Phi_d \sim d/\ln d$  is the unique positive root of the equation  $x^{d+1} = (x+1)^d$ . This almost closes the huge gap between  $\Theta(d)$  and  $\Phi_d^{d+1}$ . Our bound extends also to network congestion games. We further show that the PoS remains exponential even for singleton games. More generally, we provide a lower bound of  $\Omega((1+1/\alpha)^d/d)$  on the PoS of  $\alpha$ -approximate Nash equilibria for singleton games. All our lower bounds hold for mixed and correlated equilibria as well. On the positive side, we give a general upper bound on the PoS of  $\alpha$ -approximate Nash equilibria, which is sensitive to the range  $W$  of the player weights and the approximation parameter  $\alpha$ . We do this by explicitly constructing a novel approximate potential function, based on Faulhaber’s formula, that generalizes Rosenthal’s potential in a continuous, analytic way. From the general theorem, we deduce two interesting corollaries. First, we derive the existence of an approximate pure Nash equilibrium with PoS at most  $(d+3)/2$ ; the equilibrium’s approximation parameter ranges from  $\Theta(1)$  to  $d+1$  in a smooth way with respect to  $W$ . Second, we show that for unweighted congestion games, the PoS of  $\alpha$ -approximate Nash equilibria is at most  $(d+1)/\alpha$ .

**Key words.** congestion games, price of stability, Nash equilibrium, approximate equilibrium, potential games

**AMS subject classifications.** 68Q99, 91A10, 91A43, 90B20, 90B18

**DOI.** 10.1137/18M1207880

**1. Introduction.** In the last 20 years, a central strand of research within algorithmic game theory has focused on understanding and quantifying the inefficiency of equilibria compared to centralized, optimal solutions. There are two standard concepts that measure this inefficiency. The Price of Anarchy (PoA) [38], which takes the worst-case perspective, compares the worst-case equilibrium with the system optimum. It is a very robust measure of performance. On the other hand, the Price of Stability (PoS) [51, 5], which is also the focus of this work, takes an optimistic perspective and uses the best-case equilibrium for this comparison. The PoS is an appropriate concept for analyzing the ideal solution that we would like our protocols to produce.

The initial set of problems that arose from the PoA theory have now been resolved. The most rich and well-studied among these models are, arguably, the atomic and nonatomic variants of congestion games (see [44, Ch. 18] for a detailed discus-

---

\*Received by the editors August 16, 2018; accepted for publication (in revised form) August 23, 2019; published electronically October 24, 2019. A preliminary version of this paper appeared in ICALP’18 [21].

<https://doi.org/10.1137/18M1207880>

**Funding:** This work was supported by the Alexander von Humboldt Foundation with funds from the German Federal Ministry of Education and Research (BMBF) and by EPSRC grants EP/M008118/1 and EP/L011018/1.

<sup>†</sup>Department of Computer Science, University of Liverpool, Liverpool L69 3BX, United Kingdom (G.Christodoulou@liverpool.ac.uk, gairing@liverpool.ac.uk).

<sup>‡</sup>Chair of Operations Research, TU Munich, Munich, Bayern 80333, Germany (yian-nis.giannakopoulos@tum.de).

<sup>§</sup>Department of Computer Science, University of Liverpool, Liverpool L69 3BX, United Kingdom, and Computer Engineering and Informatics Department, University of Patras, Patras 26504, Greece (P.Spirakis@liverpool.ac.uk).

sion). This class of games is very descriptive and captures a large variety of scenarios where users compete for resources, most prominently routing games. The seminal work of Roughgarden and Tardos [49, 50] gave the answer for the nonatomic variant, where each player controls a negligible amount of traffic. Awerbuch, Azar, and Epstein [6] and Christodoulou and Koutsoupias [18] resolved the PoA for atomic congestion games with affine latencies, generalized by Aland et al. [3] to polynomials; this led to the development of Roughgarden’s smoothness framework [48], which extended the bounds to general cost functions, but also distilled and formulated previous ideas to bound the PoA in an elegant, unified framework. At the computational complexity front, we know that even for simple congestion games, finding a (pure) Nash equilibrium is a complete problem for the complexity class PLS (Polynomial Local Search) [24, 2].

Allowing the players to have different loads gives rise to the class of *weighted* congestion games [47]; this is a natural and very important generalization of congestion games, with numerous applications in routing and scheduling. Unfortunately though, an immediate dichotomy between weighted and unweighted congestion games occurs: the former may *not* even have pure Nash equilibria [41, 28, 30, 33]; as a matter of fact, it is a strongly NP-hard problem to even determine if that is the case [23]. Moreover, in such games there does not, in general, exist a potential function [43, 34], which is the main tool for proving equilibrium existence in the unweighted case.

As a result, a sharp contrast with respect to our understanding of the two aforementioned inefficiency notions arises. The PoA has been studied in depth, and general techniques for providing tight bounds are known. Moreover, the asymptotic behavior of weighted and unweighted congestion games with respect to the PoA is identical; it is  $\Theta(d/\log d)^d$  for both classes when latencies are polynomials of degree at most  $d$  [3].

The situation for the PoS, though, is completely different. For unweighted games we have a good understanding,<sup>1</sup> and the values are much lower than the PoA values, and also *tight*—approximately 1.577 for affine functions [19, 13] and  $\Theta(d)$  [17] for polynomials. For weighted games, though, there is a huge gap; the current state-of-the-art lower bound is  $\Theta(d)$ , and the upper bound is  $\Theta(d/\ln d)^d$ . These previous results are summarized at the left of Table 1.1a.

The main focus of this work is precisely to deal with this lack of understanding and to determine the PoS of weighted congestion games. What makes this problem challenging is that the only general known technique for showing upper bounds for the PoS is the potential method, which is applicable only to potential games. In a nutshell, the idea of this method is to use the global minimizer of Rosenthal’s potential [46] as an equilibrium refinement. This equilibrium is also a pure Nash equilibrium and can serve as an upper bound of the PoS. Interestingly, it turns out that, for several classes of potential games, this technique actually provides the tight answer (see, for example, [5, 19, 13, 17]). However, as already mentioned above, unlike their

---

<sup>1</sup>Much work has been also done on the PoS for *network design games*, which is, though, not so closely related to ours; in such games the cost of using an edge is split equally among players, and thus cost functions are decreasing, as opposed to our model of congestion games with nondecreasing latencies. This problem was first studied by Anshelevich et al. [5], who showed a tight bound of  $H_n$ , the harmonic number of the number of players  $n$ , for directed networks. Finding tight bounds on undirected networks is still a long-standing open problem (see, e.g., [27, 10, 39]). Recently, Bilò, Flammini, and Moscardelli [12] (asymptotically) resolved the question for broadcast networks. For the weighted variant of this problem, Albers [4] showed a lower bound of  $\Omega(\log W/\log \log W)$ , where  $W$  is the sum of the players’ weights, while Chen and Roughgarden [16] showed an upper bound of  $O(\log W/\alpha)$  for  $\alpha$ -approximate equilibria (the latter is similar in spirit to our results in section 4). See [12, 4] and references therein for a thorough discussion of those results.

TABLE 1.1

The PoA and PoS for unweighted and weighted congestion games, with polynomial latency functions of maximum degree  $d$ .  $\Phi_d$  is the unique positive solution of  $(x+1)^d = x^{d+1}$  and  $\Theta_d = \Theta(d/\log d)$ . Tight answers were known for all settings, except for the PoS of the weighted case where only trivial bounds existed. In this paper we almost close this gap by showing a lower bound of  $\Omega(\Phi_d)^{d+1}$  (Theorem 3.1), which remains exponential even for singleton games (Theorem 3.5).

(a) Previous results		(b) This paper	
	PoA	PoS	
unweighted	$\lfloor \Phi_d \rfloor^{d+1}$ [3]	$\Theta(d)$ [17]	general
weighted	$\Phi_d^{d+1}$ [3]	$[\Theta(d), \Phi_d^{d+1}]$	PoS lower bound
			$\Omega(\Phi_d)^{d+1}$
			singleton
			$\Omega(2^d/d)$
		$\alpha$ -approximate equilibria	$\Omega((1+1/\alpha)^d/d)$

unweighted counterparts, weighted congestion games are not potential games;<sup>2</sup> so, a completely fresh approach is required. One way to override the aforementioned limitations of nonexistence of pure Nash equilibria, but also their computational hardness, is to consider *approximate* equilibria. In this direction, Hansknecht, Klimm, and Skopalik [32] have shown that  $(d+1)$ -approximate pure Nash equilibria always exist in weighted congestion games with polynomial latencies of maximum degree  $d$ , while, on the negative side, there exist games that do not have 1.153-approximate pure Nash equilibria. Notice here that these results do not take into account computational complexity considerations; if we insist on polynomial-time algorithms for actually finding those equilibria, then the currently best approximation parameter becomes  $d^{O(d)}$  [14, 15, 26].

**1.1. Our results.** We provide lower and upper bounds on the PoS for the class of weighted congestion games with polynomial latencies with nonnegative coefficients. We consider both exact and approximate equilibria. Our lower bounds are summarized in Table 1.1b.

*Lower bound for weighted congestion games.* In our main result in Theorem 3.1, we resolve a long-standing open problem by providing almost tight bounds for the PoS of weighted congestion games with polynomial latency functions. We construct an instance having a PoS of  $\Omega(\Phi_d)^{d+1}$ , where  $d$  is the maximum degree of the latencies and  $\Phi_d \sim \frac{d}{\ln d}$  is the unique positive solution of equation  $(x+1)^d = x^{d+1}$ .

This bound almost closes the previously huge gap between  $\Theta(d)$  and  $\Phi_d^{d+1}$  for the PoS of weighted congestion games. The previously best lower and upper bounds were rather trivial: the lower bound corresponds to the PoS results of Christodoulou and Gairing [17] for the unweighted case (and thus, it is also a valid lower bound for the general weighted case as well), and the upper bound comes from the PoA results of Aland et al. [3] (PoA, by definition, upper-bounds PoS). It is important to make clear here that our lower bound still leaves an open gap for future work: the constant within the base of  $\Omega(\Phi_d)$  in Theorem 3.1 produces a lower bound of  $(\frac{1}{2}\Phi_d)^{d+1}$ , which is formally a factor of  $2^{d+1}$  away from the  $\Phi_d^{d+1}$  PoA upper bound.

Although, as mentioned before, weighted congestion games do not always possess pure equilibria, our lower bound construction involves a *unique* equilibrium occurring by iteratively eliminating strongly dominated strategies. As a result, this lower bound

<sup>2</sup>For the special case of weighted congestion games with linear latency functions, a potential does exist [28], and this was used by [9] to provide a PoS upper bound of 2.

holds not only for pure but also for mixed and correlated equilibria.

*Singleton games.* Next we switch to the class of singleton congestion games, where a pure strategy for each player is a *single* resource. This class is very well studied as, on one hand, it abstracts scheduling environments, and, on the other, it has very attractive equilibrium properties; unlike general weighted congestion games, there exists an (ordinal) lexicographic potential [29, 35], thus implying the existence of *pure* Nash equilibria. It is important to note that the tight lower bounds for the PoA of general weighted congestion games hold also for the class of singleton games [13, 8, 11].

Even for this special class, we show in Theorem 3.5 an exponential lower bound of  $\Omega(2^d/d)$ . The previous best lower and upper bounds were the same as those of the general case, namely  $\Theta(d)$  and  $\Phi_d^{d+1}$ , respectively. As a matter of fact, this new lower bound comes as a corollary of a more general result that we show in Theorem 3.5 that extends to approximate equilibria and gives a lower bound of  $\Omega((1+1/\alpha)^d/d)$  on the PoS of  $\alpha$ -approximate equilibria for any (multiplicative) approximation parameter  $\alpha \in [1, d]$ . Setting  $\alpha = 1$ , we recover the special case of exact equilibria and the aforementioned exponential lower bound on the standard, exact notion of the PoS. Notice here that, as we show in Theorem D.1, the optimal solution (which, in general, is not an equilibrium) itself constitutes a  $(d+1)$ -approximate equilibrium with a (trivially) optimal PoS of 1.

*Positive results for approximate equilibria.* In light of the above results, in section 4, we turn our attention to identifying environments with more structure or flexibility with respect to the underlying solution concept, for which we can hope for improved quality of equilibria. Both our lower bound constructions discussed above use players' weights that form a geometric sequence. In particular, the ratio  $W$  of the largest over the smallest weight is equal to  $w^n$  (for some  $w > 1$ ), which grows very large as the number of players  $n \rightarrow \infty$ . On the other hand, for games where the players have equal weights, i.e.,  $W = 1$ , we know that the PoS is at most  $d+1$ . It is therefore natural to ask how the performance of the good equilibria captured by the notion of PoS varies with respect to  $W$ . In Theorem 4.4, we are able to give a general upper bound for  $\alpha$ -approximate equilibria which is sensitive to this parameter  $W$  and to  $\alpha$ . This general theorem has two immediate, interesting corollaries.

First (Corollary 4.5), by allowing the ratio  $W$  to range in  $[1, \infty)$ , we derive the existence of an  $\alpha$ -approximate pure Nash equilibrium with PoS at most  $(d+3)/2$ ; the equilibrium's approximation parameter  $\alpha$  ranges from  $\Theta(1)$  to  $d+1$  in a smooth way with respect to  $W$ . This is of particular importance in settings where player weights are not very far away from each other (that is,  $W$  is small). Second (Corollary 4.6), by setting  $W = 1$  and allowing  $\alpha$  to range up to  $d+1$ , we get an upper bound of  $\frac{d+1}{\alpha}$  for the  $\alpha$ -approximate PoS of *unweighted* congestion games which, to the best of our knowledge, was not known before, degrading gracefully from  $d+1$  (which is the actual PoS of exact equilibria in the unweighted case [17]) down to the optimal value of 1 if we allow  $(d+1)$ -approximate equilibria (which in fact can be achieved by the optimum solution itself; see Theorem D.1).

*Our techniques.* An advantage of our main lower bound (Theorem 3.1) is the simplicity of the underlying construction as well as its straightforward adaptation to network games (see subsection 3.1.1). However, fine-tuning the parameters of the game (player weights and latency functions), to ensure uniqueness of the equilibrium at the "bad" instance, was a technically involved task. This was due in part to the fact that, in order to guarantee uniqueness (via iteratively dominant strategies), each player interacts with a window of  $\mu$  other players. This  $\mu$  depends on  $d$  in a delicate way (see Figure 3.1 and Lemma 3.2); it has to be an integer but, at the same time,

needs also to balance nicely with the algebraic properties of  $\Phi_d$ .

Moreover, we needed to provide deeper insights into the asymptotic, analytic behavior of  $\Phi_d$  and to explore some new algebraic characteristics of  $\Phi_d$  (see, e.g., Lemma A.3). It is important to keep in mind that asymptotically  $\Phi_d \sim \frac{d}{\ln d}$  (see (A.6)). The fact that  $\Phi_d = \Theta\left(\frac{d}{\ln d}\right)$  was already known by the work of Aland et al. [3]; here we provide a more refined characterization of  $\Phi_d$ 's growth, using the Lambert-W function (see (A.7)).

In order to derive our upper bounds, we need to define a novel *approximate potential function* [16, 20, 32]. First, in Lemma 4.1, we identify clear algebraic sufficient conditions for the existence of approximate equilibria with good social-cost guarantees, and then we explicitly define (see (4.4) and (4.10) in the proof of Theorem 4.4) a function that satisfies them. This continuous function, which is defined in the entire space of positive reals, essentially generalizes that of Rosenthal in a smooth way: by setting  $W = \alpha = 1$ , we recover exactly the first significant terms of the well-known Rosenthal potential [46] polynomial, with which one can demonstrate the usual PoS results for the unweighted case (see, e.g., [19]). The simple, analytic way in which this function is defined is the very reason we can handle both the approximation parameter  $\alpha$  of the equilibrium and the ratio  $W$  of the weights in a smooth manner while at the same time providing good PoS guarantees.

It is important to stress that, by the purely analytical way in which our approximate potential function is defined, in principle it can also incorporate more general cost functions than polynomials; so, we believe that this technique may be of independent interest. We point towards that direction in Appendix C.

**2. Model and notation.** Let  $\mathbb{R}$  denote the set of real numbers,  $\mathbb{R}_{\geq 0} = [0, \infty)$  and  $\mathbb{R}_{> 0} = (0, \infty)$ .

*Weighted congestion games.* A *weighted congestion game* consists of a finite, nonempty set of players  $N$  and resources (or facilities)  $E$ . Each player  $i \in N$  has a *weight*  $w_i \in \mathbb{R}_{> 0}$  and a *strategy set*  $S_i \subseteq 2^E$ . Associated with each resource  $e \in E$  is a *cost* (or *latency*) *function*  $c_e : \mathbb{R}_{> 0} \rightarrow \mathbb{R}_{\geq 0}$ . In this paper we mainly focus on polynomial cost functions with maximum degree  $d \geq 0$  and nonnegative coefficients; that is, every cost function is of the form  $c_e(x) = \sum_{j=0}^d a_{e,j} \cdot x^j$ , with  $a_{e,j} \geq 0$  for all  $j$ . In the following, whenever we refer to polynomial cost functions we mean cost functions of this particular form.

A *pure strategy profile* is a choice of strategies  $\mathbf{s} = (s_1, s_2, \dots, s_n) \in S = S_1 \times \dots \times S_n$  by the players. We use the standard game-theoretic notation  $\mathbf{s}_{-i} = (s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_n)$ ,  $S_{-i} = S_1 \times \dots \times S_{i-1} \times S_{i+1} \times \dots \times S_n$ , such that  $\mathbf{s} = (s_i, \mathbf{s}_{-i})$ . Given a pure strategy profile  $\mathbf{s}$ , we define the *load*  $x_e(\mathbf{s})$  of resource  $e \in E$  as the total weight of players that use resource  $e$  on  $\mathbf{s}$ , i.e.,  $x_e(\mathbf{s}) = \sum_{i \in N: e \in s_i} w_i$ . The *cost* player  $i$  is defined by  $C_i(\mathbf{s}) = \sum_{e \in s_i} c_e(x_e(\mathbf{s}))$ .

A *singleton* weighted congestion game is a special form of congestion game where the strategies of all players consist only of single resources; that is, for all players  $i \in N$ ,  $|s_i| = 1$  for all  $s_i \in S_i$ . In a weighted *network* congestion game the resources  $E$  are given as the edge set of some directed graph  $G = (V, E)$ , and each player  $i \in N$  has a source  $o_i \in V$  and destination  $t_i \in V$  node; then, the strategy set  $S_i$  of each player is implicitly given as the edge sets of all directed  $o_i \rightarrow t_i$  paths in  $G$ .

*Nash equilibria.* A pure strategy profile  $\mathbf{s}$  is a pure *Nash equilibrium* if and only if for every player  $i \in N$  and for all  $s'_i \in S_i$ , we have  $C_i(\mathbf{s}) \leq C_i(s'_i, \mathbf{s}_{-i})$ . Similarly a strategy profile is an  $\alpha$ -*approximate pure Nash equilibrium*, for  $\alpha \geq 1$ , if  $C_i(\mathbf{s}) \leq \alpha \cdot C_i(s'_i, \mathbf{s}_{-i})$  for all players  $i \in N$  and  $s'_i \in S_i$ . As discussed in the introduction, weighted

congestion games do not always admit pure Nash equilibria. However, by Nash’s theorem they have mixed Nash equilibria. A tuple  $\sigma = (\sigma_1, \dots, \sigma_N)$  of independent probability distributions over players’ strategy sets is a *mixed Nash equilibrium* if

$$\mathbb{E}_{\mathbf{s} \sim \sigma} [C_i(\mathbf{s})] \leq \mathbb{E}_{\mathbf{s}_{-i} \sim \sigma_{-i}} [C_i(s'_i, \mathbf{s}_{-i})]$$

holds for every  $i \in N$  and  $s'_i \in S_i$ . Here  $\sigma_{-i}$  is a product distribution of all  $\sigma_j$ ’s with  $j \neq i$ , and  $\mathbf{s}_{-i}$  denotes a strategy profile drawn from this distribution. We use  $\text{NE}(G)$  to denote the set of all mixed Nash equilibria of a game  $G$ .

*Social cost and price of stability.* Fix a weighted congestion game  $G$ . The *social cost* of a pure strategy profile  $\mathbf{s}$  is the weighted sum of the players’ costs

$$C(\mathbf{s}) = \sum_{i \in N} w_i \cdot C_i(\mathbf{s}) = \sum_{e \in E} x_e(\mathbf{s}) \cdot c_e(x_e(\mathbf{s})).$$

Denote by  $\text{OPT}(G) = \min_{\mathbf{s} \in S} C(\mathbf{s})$  the *optimum social cost* over all strategy profiles  $\mathbf{s} \in S$ . Then, the *Price of Stability (PoS)* of  $G$  is the social cost of the best-case Nash equilibrium over the optimum social cost:

$$\text{PoS}(G) = \min_{\sigma \in \text{NE}(G)} \frac{\mathbb{E}_{\mathbf{s} \sim \sigma} [C(\mathbf{s})]}{\text{OPT}(G)}.$$

The PoS of  $\alpha$ -approximate Nash equilibria is defined accordingly. The PoS for a class  $\mathcal{G}$  of games is the worst (i.e., largest) PoS among all games in the class, that is,  $\text{PoS}(\mathcal{G}) = \sup_{G \in \mathcal{G}} \text{PoS}(G)$ . For example, our focus in this paper is determining the PoS for the class  $\mathcal{G}$  of weighted congestion games with polynomial cost functions.

For brevity, we will sometimes abuse our formal terminology and refer to the “PoS of  $\mathbf{s}$ ” for a specific (approximate) equilibrium  $\mathbf{s}$  of a game  $G$  (see, e.g., Theorem 4.4); by that we will mean the approximation ratio of the social cost of  $\mathbf{s}$  to the optimum, i.e.,  $\frac{C(\mathbf{s})}{\text{OPT}(G)}$ . Clearly, the PoS of *any* such equilibrium  $\mathbf{s}$  is a valid upper bound to the PoS of the entire game  $G$ .

Finally, notice that, by using a straightforward scaling argument, it is without loss with respect to the PoS metric to analyze games with player weights in  $[1, \infty)$ ; if not, divide all  $w_i$ ’s with  $\min_i w_i$  and scale cost functions accordingly.

**3. Lower bounds.** In this section, we present our lower bound constructions. In subsection 3.1 we present the general lower bound and then in subsection 3.2 the lower bound for singleton games.

**3.1. General congestion games.** The next theorem presents our main negative result on the PoS of weighted congestion games with polynomial latencies of degree  $d$  that almost matches the PoA upper bound of  $\Phi_d^{d+1}$  from Aland et al. [3]. Our result shows a strong separation for the PoS between weighted and unweighted congestion games; the PoS of the latter is at most  $d + 1$  [17]. This is in sharp contrast to the PoA of these two classes, where the respective bounds are essentially the same.

To state our result, we first introduce some notation. Let  $\Phi_d \sim \frac{d}{\ln d}$  be the unique positive root of the equation  $(x+1)^d = x^{d+1}$ , and let  $\beta_d$  be a parameter with  $\beta_d \geq 0.38$  for any  $d$ ,  $\lim_{d \rightarrow \infty} \beta_d = \frac{1}{2}$  (formally,  $\beta_d$  is defined in (3.2) below, and a plot of its values can be seen in Figure 3.1).

**THEOREM 3.1.** *The PoS of weighted congestion games with polynomial latency functions of degree at most  $d \geq 9$  is at least  $(\beta_d \Phi_d)^{d+1}$ .*




FIG. 3.1. The values of parameters  $\beta_b$  and  $c_d$  in Lemma 3.2 and Theorem 3.1 for  $d = 9, 10, \dots, 100$ .

We must mention here that the restriction of  $d \geq 9$  is without loss: for polynomial latencies of smaller degrees  $d \leq 8$  we can instead apply the simpler lower bound instance for singleton games given in subsection 3.2. To prove the theorem, we will need the following technical lemma. Its proof can be found in Appendix A.2.

LEMMA 3.2. For any positive integer  $d$  define

$$(3.1) \quad c_d = \frac{1}{d} \left\lceil d \frac{\ln(2 \cdot \Phi_d + 1) - \ln(\Phi_d + 1)}{\ln \Phi_d} \right\rceil$$

and

$$(3.2) \quad \beta_d = 1 - \Phi_d^{-c_d}.$$

Then

$$(3.3) \quad \Phi_d^{d+2} \leq \left( \Phi_d + \frac{1}{\beta_d} \right)^d,$$

and for all  $d \geq 9$ ,

$$(3.4) \quad d \cdot c_d \geq 3, \quad 0.38 \leq \beta_d \leq \frac{1}{2}, \quad \text{and} \quad \lim_{d \rightarrow \infty} \beta_d = \frac{1}{2}.$$

Plots of parameters  $c_d$  and  $\beta_d$  can be found in Figure 3.1.

*Proof of Theorem 3.1.* Fix some integer  $d \geq 9$ . Our lower bound instance consists of  $n + \mu$  players and  $n + \mu + 1$  facilities, where  $\mu := c \cdot d$  for  $c = c_d$  defined as in (3.1). In particular then, due to (3.4) of Lemma 3.2,  $\mu \geq 3$  is an integer. We can think of  $n$  as a very large integer, since at the end we will take  $n \rightarrow \infty$ . Every player  $i = 1, 2, \dots, n + \mu$  has a weight of  $w_i = w^i$ , where  $w = 1 + \frac{1}{\Phi_d}$ .

It will be useful for subsequent computations to notice that

$$w^d = \left(1 + \frac{1}{\Phi_d}\right)^d = \frac{(\Phi_d + 1)^d}{\Phi_d^d} = \frac{\Phi_d^{d+1}}{\Phi_d^d} = \Phi_d,$$

$$w^{d+1} = w^d \cdot w = \Phi_d \left(1 + \frac{1}{\Phi_d}\right) = \Phi_d + 1.$$

Let us also define

$$\alpha = \alpha(\mu) := \sum_{j=1}^{\mu} w^{-j} = \frac{1 - w^{-\mu}}{w - 1} = \frac{1 - (w^d)^{-c}}{w - 1} = \frac{1 - \Phi_d^{-c}}{1 + \frac{1}{\Phi_d} - 1} = \Phi_d (1 - \Phi_d^{-c}) = \beta \Phi_d,$$

where  $\beta = \beta_d$  is defined as in (3.2). In the following we will make extensive use of the observation that

$$w^{-\mu} = (w^d)^{-c} = \Phi_d^{-c} = 1 - \beta.$$

Furthermore, for every  $i \geq \mu + 1$

$$\sum_{j=i-\mu}^{i-1} w_j = \sum_{j=1}^{\mu} w^{i-j} = \alpha \cdot w^i \quad \text{and} \quad \sum_{j=i-\mu}^i w_j = (\alpha + 1) \cdot w^i,$$

and

$$\sum_{\ell=1}^{\infty} w^{-\ell} = \frac{1}{w - 1} = \frac{1}{1 + \frac{1}{\Phi_d} - 1} = \Phi_d.$$

The facilities have latency functions

$$\begin{aligned} c_j(t) &= \Phi_d(1 - \beta)(\alpha + 1)^d && \text{if } j = 1, \dots, \mu, \\ c_j(t) &= w^{-j(d+1)}t^d && \text{if } j = \mu + 1, \dots, \mu + n, \\ c_{n+\mu+1}(t) &= 0. \end{aligned}$$

Every player  $i$  has two available strategies,  $s_i^*$  and  $\tilde{s}_i$ . Eventually we will show that the profile  $\mathbf{s}^*$  corresponds to the optimal solution, while  $\tilde{\mathbf{s}}$  corresponds to the *unique* Nash equilibrium of the game. Informally, at the former the player chooses to stay at her “own”  $i$ th facility, while at the latter she chooses to deviate and play the  $\mu$  following facilities  $i + 1, \dots, i + \mu$ . However, special care shall be taken for the boundary cases of the first  $\mu$  and last  $\mu$  players, so for any player  $i$  we formally define  $S_i = \{s_i^*, \tilde{s}_i\}$ , where  $s_i^* = \{i\}$  and

$$\tilde{s}_i = \begin{cases} \{\mu + 1, \dots, \mu + i\} & \text{if } i = 1, \dots, \mu, \\ \{i + 1, \dots, i + \mu\} & \text{if } i = \mu + 1, \dots, n, \\ \{i + 1, \dots, n + \mu + 1\} & \text{if } i = n + 1, \dots, n + \mu. \end{cases}$$

These two outcomes,  $\mathbf{s}^*$  and  $\tilde{\mathbf{s}}$ , are shown in Figure 3.2.

Notice here that any facility  $j$  cannot get a load greater than the sum of the weights of the previous  $\mu$  players plus the weight of the  $j$ th player. So, for any strategy profile  $\mathbf{s}$ ,

$$(3.5) \quad x_j(\mathbf{s}) \leq \sum_{\ell=j-\mu}^j w_\ell = (\alpha + 1)w^j \quad \text{for all } j \geq \mu + 1.$$

Next we will show that the strategy profile  $\tilde{\mathbf{s}} = (\tilde{s}_1, \dots, \tilde{s}_{n+\mu})$  is the *unique* Nash equilibrium of our congestion game. We do that by proving the following:






FIG. 3.2. The social optimum  $\mathbf{s}^*$  and the unique Nash equilibrium  $\tilde{\mathbf{s}}$  in the lower bound construction of Theorem 3.1 for general weighted congestion games.

1. It is a strongly *dominant* strategy for any player  $i = 1, \dots, \mu$  to play  $\tilde{s}_i$ .
2. For any  $i = \mu + 1, \dots, n + \mu$ , given that every player  $k < i$  has chosen to play  $\tilde{s}_k$ , then it is a strongly *dominant* strategy for player  $i$  to deviate to  $\tilde{s}_i$  as well.

For the first condition, fix some player  $i \leq \mu$  and a strategy profile  $\mathbf{s}_{-i}$  for the other players, and observe that by choosing  $\tilde{s}_i$ , player  $i$  incurs a cost of at most

$$\begin{aligned}
 C_i(\tilde{s}_i, \mathbf{s}_{-i}) &= \sum_{j \in \tilde{s}_i} c_j(x_j(\tilde{s}_i)) \leq \sum_{\ell=\mu+1}^{\mu+i} c_\ell((\alpha + 1)w^\ell) \\
 &= \sum_{\ell=\mu+1}^{\mu+i} w^{-\ell(d+1)}(\alpha + 1)^d w^{\ell d} = (\alpha + 1)^d \sum_{\ell=\mu+1}^{\mu+i} w^{-\ell} \\
 &< (\alpha + 1)^d \cdot w^{-\mu} \cdot \sum_{\ell=1}^{\infty} w^{-\ell} = (\alpha + 1)^d \cdot (1 - \beta) \cdot \Phi_d \\
 &= C_i(s_i^*, \mathbf{s}_{-i}),
 \end{aligned}$$

where in the first inequality we used the bound from (3.5).

For the second condition, we will consider the deviations of the remaining players. Fix now some  $i = \mu + 1, \dots, n$ , and assume a strategy profile  $\mathbf{s}_{-i} = (\tilde{s}_1, \dots, \tilde{s}_{i-1}, s_{i+1}, \dots, s_{n+\mu})$  for the remaining players.<sup>3</sup> If player  $i$  chooses strategy  $s_i^*$ , she will experience a cost of

$$C_i(s_i^*, \mathbf{s}_{-i}) = c_i \left( \sum_{\ell=i-\mu}^i w_\ell \right) = c_i((\alpha + 1)w^i) = w^{-i(d+1)}(\alpha + 1)^d w^{id} = (\alpha + 1)^d w^{-i}.$$

<sup>3</sup>For the remaining  $\mu$  players  $i = n + 1, \dots, n + \mu$  the proof is similar, and as a matter of fact easier, since when these players deviate to  $\tilde{s}_i$  they also use the final “dummy” facility  $n + \mu + 1$  that has zero cost.




FIG. 3.3. The format of profile  $\mathbf{s}'_{-i}$  described in Claim 3.3 and returned as output from Procedure DOMINATE( $\mathbf{s}_{-i}, i$ ) (see Appendix A.3). All players  $i + 1, \dots, i + \mu$  (i.e., those who lie within the window of interest of player  $i$ , depicted in gray) play according to the Nash equilibrium  $\tilde{\mathbf{s}}$ , except the last player  $i + \mu$  (that plays according to the optimal profile  $\mathbf{s}^*$ ) and at most one other  $k$  (that may play either  $\tilde{s}_k$  or  $s_k^*$ ).

It remains to show that

$$(3.6) \quad C_i(\tilde{\mathbf{s}}_i, \mathbf{s}_{-i}) < C_i(\mathbf{s}_i^*, \mathbf{s}_{-i}) = (\alpha + 1)^d w^{-i}.$$

The cost  $C_i(\tilde{\mathbf{s}}_i, \mathbf{s}_{-i})$  is complicated to bound immediately for any profile  $\mathbf{s}_{-i}$ . Instead, we will resort to the following claim, which characterizes the profile  $\mathbf{s}_{-i}$  where this cost is maximized, as shown in Figure 3.3. Its proof can be found in Appendix A.3.

- CLAIM 3.3. *There exists a profile  $\mathbf{s}'_{-i}$  such that*
1.  $s'_j = s_j$  for all  $j < i$  and  $j > i + \mu$ ;
  2.  $s'_{i+\mu} = s_{i+\mu}^*$ ;
  3. *there exists some  $k \in \{i + 1, \dots, i + \mu - 1\}$  such that*

$$s'_j = \tilde{s}_j \quad \text{for all } j \in \{i + 1, \dots, i + \mu - 1\} \setminus \{k\},$$

which dominates  $\mathbf{s}_{-i}$ , i.e.,

$$(3.7) \quad C_i(\tilde{\mathbf{s}}_i, \mathbf{s}_{-i}) \leq C_i(\tilde{\mathbf{s}}_i, \mathbf{s}'_{-i}).$$

By use of Claim 3.3, it remains to show that

$$(3.8) \quad C_i(\tilde{\mathbf{s}}_i, \mathbf{s}'_{-i}) < (\alpha + 1)^d w^{-i},$$

just for the special case of profiles  $\mathbf{s}'$  that are described in Claim 3.3 and also shown in Figure 3.3. We do this in Appendix A.4.

Summarizing, we proved that indeed  $\tilde{\mathbf{s}}$  is the *unique* Nash equilibrium of our congestion game. Finally, to conclude with lower-bounding the PoS, let us compute the social cost on profiles  $\tilde{\mathbf{s}}$  and  $\mathbf{s}^*$ . On  $\mathbf{s}^*$ , any facility  $j$  (except the last one) gets a

load equal to the weight of player  $j$ , so

$$\begin{aligned}
 C(\mathbf{s}^*) &= \sum_{j=1}^{n+\mu} w_j c_j(w_j) \\
 &= \sum_{j=1}^{\mu} w^j \Phi_d (1-\beta)(\alpha+1)^d + \sum_{j=\mu+1}^{n+\mu} w^j w^{-j(d+1)} (w^j)^d \\
 &= \Phi_d (1-\beta)(\alpha+1)^d \sum_{j=1}^{\mu} w^j + \sum_{j=\mu+1}^{\mu+n} 1 \\
 &= \Phi_d (1-\beta)(\alpha+1)^d w \frac{w^{\mu}-1}{w-1} + n \\
 &= \Phi_d (1-\beta)(\alpha+1)^d \left(1 + \frac{1}{\Phi_d}\right) \frac{\frac{1}{1-\beta}-1}{1 + \frac{1}{\Phi_d}-1} + n \\
 &= \Phi_d (1-\beta)(\alpha+1)^d (\Phi_d+1) \frac{\beta}{1-\beta} + n \\
 &\leq n + \Phi_d (\Phi_d+1) \beta (\alpha+1)^d.
 \end{aligned}$$

On the other hand, at the unique Nash equilibrium  $\tilde{\mathbf{s}}$  each facility  $j \geq \mu + 1$  receives a load equal to the sum of the weights of the previous  $\mu$  players, i.e.,

$$x_j(\tilde{\mathbf{s}}) = \sum_{\ell=j-\mu}^{j-1} w_{\ell} = \alpha w^j,$$

so

$$C(\tilde{\mathbf{s}}) \geq \sum_{j=\mu+1}^{n+\mu} x_j(\tilde{\mathbf{s}}) c_j(x_j(\tilde{\mathbf{s}})) = \sum_{j=\mu+1}^{n+\mu} w^{-j(d+1)} (\alpha w^j)^{d+1} = \alpha^{d+1} \sum_{j=\mu+1}^{\mu+n} 1 = \alpha^{d+1} n.$$

By taking  $n$  arbitrarily large we get a lower bound on the PoS of

$$\lim_{n \rightarrow \infty} \frac{C(\tilde{\mathbf{s}})}{C(\mathbf{s}^*)} \geq \lim_{n \rightarrow \infty} \frac{\alpha^{d+1} n}{n + \Phi_d (\Phi_d + 1) \beta (\alpha + 1)^d} = \alpha^{d+1} = (\beta \Phi_d)^{d+1},$$

where from Lemma 3.2 we know that  $\frac{1}{3} \leq \beta = \frac{1}{2} - o(1)$ . □

**3.1.1. Network games.** Due to the rather simple structure of the players’ strategy sets in the lower bound construction of Theorem 3.1, it can be readily extended to network games as well.

PROPOSITION 3.4. *Theorem 3.1 applies also to network weighted congestion games.*

*Proof.* We arrange the resources from the instance in the proof of Theorem 3.1 as edges in a graph as depicted in Figure 3.4. In particular, for all  $j = \mu + 1, \dots, n + \mu + 1$ , resource  $j$  from Theorem 3.1 corresponds to edge  $(u_j, u_{j+1})$ . For the special case of the first  $\mu$  resources, for  $j = 1, \dots, \mu$ , we represent resource  $j$  by the directed edge  $(u_{\mu+1}, u_j)$ .

Regarding strategies, recall from the instance used in the proof of Theorem 3.1 that, for each  $i = \mu, \dots, n$ , player  $i$  has two available strategies: resource  $\{i\}$  and resources  $\{i + 1, \dots, i + \mu\}$ . To map this to our network instance, we set the source




FIG. 3.4. Transformation of the lower bound instance of Theorem 3.1 for general weighted congestion games to a network game, as described in Proposition 3.4.

node  $o_i$  of player  $i$  to be  $o_i = u_{i+1}$  and introduce a new destination node  $t_i$  connected to the rest of the graph by zero-latency directed edges  $(u_i, t_i)$  and  $(u_{i+\mu+1}, t_i)$ . In that way, strategy  $\{i\}$  of Theorem 3.1 corresponds to the path  $o_i = u_{i+1} \rightarrow u_i \rightarrow t_i$  of our graph, while strategy  $\{i+1, \dots, i+\mu\}$  corresponds to path  $o_i = u_{i+1} \rightarrow u_{i+2} \rightarrow \dots \rightarrow u_{i+\mu+1} \rightarrow t_i$ . To avoid clutter, these destination nodes  $t_i$  and the corresponding zero-latency edges are not depicted in Figure 3.4. In an analogous way, we can set the sources and destinations of the remaining first  $i = 1, \dots, \mu - 1$  and last  $i = n + 1, \dots, n + \mu + 1$  players, taking into consideration their specially restricted strategy sets in the construction of the proof of Theorem 3.1.

Summarizing, each player  $i \in [1, n + \mu]$  has to route its traffic from  $o_i$  to  $t_i$ , where

$$o_i = \begin{cases} u_{\mu+1} & \text{if } i = 1, \dots, \mu, \\ u_{i+1} & \text{if } i = \mu + 1, \dots, n + \mu, \end{cases}$$

and nodes  $t_i$  are connected with *zero-latency* edges as follows:

- For each  $i \in [1, n + \mu]$  there is a directed zero cost edge from  $u_i$  to  $t_i$ .
- For each  $i \in [1, n]$  there is a directed zero cost edge from  $u_{\mu+1+i}$  to  $t_i$ .
- For each  $i \in [n + 1, n + \mu]$  there is a directed zero cost edge from  $u_{\mu+n+2}$  to  $t_i$ .

Then, by construction, each player  $i$  has two available  $o_i \rightarrow t_i$  paths, which correspond directly to strategy sets  $s_i^*$  and  $\tilde{s}_i$  used in the proof of Theorem 3.1.

There is one issue left to complete our network game construction: we have not yet set a *direction* on some of our edges, namely  $(u_i, u_{i+1})$  for  $i = \mu + 1, \dots, n + \mu + 1$ , which are depicted also as undirected edges in Figure 3.4. This is due to the fact that, by our construction so far, these edges can be used in *both* directions: by player  $i$  (left-to-right) or player  $i + 1$  (right-to-left). Thus, to turn our instance to a valid *directed* network, we need to replace such edges with a “gadget” that essentially forces both players, no matter from which direction they enter the edge, to use it in the same direction and *both* contribute to its load. This can be achieved by using the structure depicted in the bottom right corner of Figure 3.4.  $\square$

**3.2. Singleton games.** In this section we give an exponential lower bound for *singleton* weighted congestion games with polynomial latency functions. The following theorem also handles approximate equilibria and provides a lower bound on the PoS in a very strong sense; even if one allows for the best approximate equilibrium with approximation factor  $\alpha = o\left(\frac{d}{\ln d}\right)$ , then its cost is lower-bounded by  $\omega(\text{poly}(d))$  times

the optimal cost.<sup>4</sup> In other words, in order to achieve polynomial (with respect to  $d$ ) guarantees on the PoS, one has to consider  $\Omega\left(\frac{d}{\ln d}\right)$ -approximate equilibria—almost linear in  $d$ ; this shows that our positive result in Corollary 4.5, of the following subsection 4.3, is almost tight. This is furthermore complemented by Theorem D.1, where we show that the socially optimum profile is a  $(d + 1)$ -approximate equilibrium (achieving an optimal PoS of 1).

**THEOREM 3.5.** *For any positive integer  $d$  and any real  $\alpha \in [1, d)$ , the  $\alpha$ -approximate (mixed) PoS of weighted (singleton) congestion games with polynomial latencies of degree at most  $d$  is at least*

$$(3.9) \quad \frac{1}{e(d+1)} \left(1 + \frac{1}{\alpha}\right)^{d+1}.$$

*In particular, for the special case of  $\alpha = 1$ , we derive that the PoS of exact equilibria is  $\Omega(2^d/d)$ .*

*Proof.* Fix a positive integer  $d$  and the desired approximation parameter  $\alpha \in [1, d)$ . Also, let  $\gamma \in (\alpha, d)$  be a parameter arbitrarily close to  $\alpha$ . Our instance consists of  $n$  players with weights  $w_i = w^i, i = 1, 2, \dots, n$ , where we set

$$(3.10) \quad w = \gamma \frac{d+1}{d-\gamma} > \gamma,$$

the inequality holding due to the fact that  $d + 1 > d - \gamma > 0$ . At the end of our construction we will take  $n \rightarrow \infty$ , so one can think of  $n$  as a very large integer. There are  $n + 1$  facilities with latency functions

$$\begin{aligned} c_1(t) &= \gamma w^d (w + 1)^d, \\ c_j(t) &= (\gamma w^d)^{2-j} \cdot t^d, & j = 2, \dots, n, \\ c_{n+1}(t) &= \gamma^{1-n} w^d (w + 1)^d. \end{aligned}$$

Any player  $i$  has exactly two strategies,  $s_i^* = \{i\}$  and  $\tilde{s}_i = \{i + 1\}$ , i.e.,  $S_i = \{\{i\}, \{i + 1\}\}$  for all  $i = 1, \dots, n$ . Let  $\mathbf{s}^*, \tilde{\mathbf{s}}$  be the strategy profiles where every player  $i$  plays  $s_i^*, \tilde{s}_i$ , respectively. These two outcomes,  $\mathbf{s}^*$  and  $\tilde{\mathbf{s}}$ , are depicted in Figure 3.5. One should think of  $\mathbf{s}^*$  as the socially optimal profile. We will show that  $\tilde{\mathbf{s}}$  is the *unique*  $\alpha$ -approximate Nash equilibrium of our game. To ensure this, it suffices to require the following, which corresponds to eliminating all other possible strictly dominated  $\alpha$ -approximate equilibria:

1. It is a strictly  $\alpha$ -dominant strategy for player 1 to use facility 2, i.e.,

$$\alpha C_1(\tilde{s}_1, \mathbf{s}_{-1}) < C_1(\mathbf{s})$$

for any profile  $\mathbf{s}$ .

2. For any  $i = 2, \dots, n$ , if every player  $k < i$  has chosen facility  $k + 1$ , then it is a strictly  $\alpha$ -dominant strategy for player  $i$  to choose facility  $i + 1$ , i.e.,

$$\alpha C_i(\tilde{s}_1, \dots, \tilde{s}_{i-1}, \tilde{s}_i, s_{i+1}, \dots, s_n) < C_i(\tilde{s}_1, \dots, \tilde{s}_{i-1}, s_i, s_{i+1}, \dots, s_n)$$

for any strategies  $(s_i, s_{i+1}, \dots, s_n) \in S_i \times \dots \times S_n$ .

---

<sup>4</sup>To see this, just take any upper bound of  $\frac{d+1}{c \ln(d+1)}$  on  $\alpha$  for a constant  $c > 2$ . Then, the lower bound in (3.9) becomes  $\Omega(d^{c-1})$ .




FIG. 3.5. The social optimum  $\mathbf{s}^*$  and the unique  $\alpha$ -approximate equilibrium  $\tilde{\mathbf{s}}$  in the lower bound construction of Theorem 3.5 for singleton weighted congestion games.

For the first condition, since facility 2 can be used by at most players 1 and 2, and  $\gamma > \alpha$ , it is enough to show that  $\gamma c_2(w_1 + w_2) \leq c_1(w_1)$ . Indeed,

$$\gamma c_2(w_1 + w_2) = \gamma(\gamma w^d)^{2-2}(w + w^2)^d = \gamma w^d(1 + w)^d = c_1(w_1).$$

Similarly, for the second condition, it suffices to show that  $\gamma c_{i+1}(w_i + w_{i+1}) \leq c_i(w_{i-1} + w_i)$  for  $i = 2, \dots, n - 1$ , and  $\gamma c_{n+1}(w_n) \leq c_n(w_{n-1} + w_n)$  for the special case of  $i = n$ . This is because facility  $i + 1$  can be used by at most players  $i$  and  $i + 1$ , while facility  $i$  is already being used by player  $i - 1$ . Indeed, for any  $i = 2, \dots, n - 1$  we see that

$$\begin{aligned} \gamma c_{i+1}(w_i + w_{i+1}) &= \gamma(\gamma w^d)^{2-(i+1)}(w^i + w^{i+1})^d = (\gamma w^d)^{2-i}(w^{i-1} + w^i)^d \\ &= c_i(w_{i-1} + w_i), \end{aligned}$$

while, for  $i = n$ ,

$$\begin{aligned} c_n(w_{n-1} + w_n) &= (\gamma w^d)^{2-n}(w^{n-1} + w^n)^d \\ &= \gamma^{2-n} w^{d(2-n)+d(n-1)}(w + 1)^d \\ &= \gamma \cdot \gamma^{1-n} w^d (w + 1)^d \\ &= \gamma c_{n+1}(w_n). \end{aligned}$$

The social cost at equilibrium  $\tilde{\mathbf{s}}$  is at least the cost of player  $n$  at  $\tilde{\mathbf{s}}$ , that is,

$$C(\tilde{\mathbf{s}}) \geq w_n c_{n+1}(w_n) = w^n \cdot \gamma^{1-n} w^d (1 + w)^d = \left(\frac{w}{\gamma}\right)^n \gamma \cdot w^d (1 + w)^d.$$

On the other hand, consider the strategy profile  $\mathbf{s}^*$  where every player  $i$  chooses facility

$i$ :

$$\begin{aligned}
 C(\mathbf{s}^*) &= w_1 c_1(w_1) + \sum_{i=2}^n w_i c_i(w_i) \\
 &= \gamma w^{d+1} (1+w)^d + \sum_{i=2}^n w^i (\gamma w^d)^{2-i} w^{id} \\
 &= \gamma w^{d+1} (1+w)^d + \gamma^2 w^{2d} \sum_{i=2}^n \left(\frac{w}{\gamma}\right)^i \\
 &= \gamma w^{d+1} (1+w)^d + \gamma^2 w^{2d} \cdot \left(\frac{w}{\gamma}\right)^2 \frac{\left(\frac{w}{\gamma}\right)^{n-1} - 1}{\frac{w}{\gamma} - 1} \\
 &\leq \gamma w^{d+1} (1+w)^d + \gamma^2 w^{2d} \cdot \left(\frac{w}{\gamma}\right)^2 \frac{\left(\frac{w}{\gamma}\right)^{n-1}}{\frac{w}{\gamma} - 1} \\
 &= \left(\frac{w}{\gamma}\right)^n \gamma \cdot \left[ \left(\frac{w}{\gamma}\right)^{-n} \cdot w^{d+1} (w+1)^d + \frac{w^{2d+1}}{\frac{w}{\gamma} - 1} \right].
 \end{aligned}$$

Recall now that, from (3.10),  $\frac{w}{\gamma} > 1$ , and thus  $\lim_{n \rightarrow \infty} \left(\frac{w}{\gamma}\right)^{-n} = 0$ . So, as the number of players  $n$  grows large, we get the following lower bound on the PoS:

$$\lim_{n \rightarrow \infty} \frac{C(\tilde{\mathbf{s}})}{C(\mathbf{s}^*)} \geq \lim_{n \rightarrow \infty} \frac{w^d (1+w)^d}{\left(\frac{w}{\gamma}\right)^{-n} \cdot w^{d+1} (w+1)^d + \frac{w^{2d+1}}{\frac{w}{\gamma} - 1}} = \left(\frac{w}{\gamma} - 1\right) \frac{(1+w)^d}{w^{d+1}}.$$

Since  $\gamma$  is chosen arbitrarily close to  $\alpha$ , deploying (3.10) to substitute  $w$ , the above lower bound can be written as

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \frac{C(\tilde{\mathbf{s}})}{C(\mathbf{s}^*)} &\geq \left(\frac{d+1}{d-\alpha} - 1\right) \left[1 + \frac{\alpha(d+1)}{d-\alpha}\right]^d / \left[\frac{\alpha(d+1)}{d-\alpha}\right]^{d+1} \\
 &= \left(\frac{\alpha+1}{d-\alpha}\right) \left[\frac{d(\alpha+1)}{d-\alpha}\right]^d / \left[\frac{\alpha(d+1)}{d-\alpha}\right]^{d+1} \\
 &= \frac{(\alpha+1)^{d+1} d^d}{\alpha^{d+1} (d+1)^{d+1}} \\
 &= \frac{1}{d+1} \left(1 - \frac{1}{d+1}\right)^d \left(1 + \frac{1}{\alpha}\right)^{d+1} \\
 &\geq \frac{1}{e} \frac{1}{d+1} \left(1 + \frac{1}{\alpha}\right)^{d+1}. \quad \square
 \end{aligned}$$

**4. Upper bounds.** The negative results of the previous sections involve constructions where the ratio  $W$  of the largest to smallest weight can be exponential in  $d$ . In the main theorem (Theorem 4.4) of this section we present an analysis which is sensitive to this parameter  $W$ , and we identify conditions under which the performance of approximate equilibria can be significantly improved.

Our upper bound approach is based on the design of a suitable approximate potential function and has three main steps. First, in subsection 4.1, we set up a framework for the definition of this function by identifying conditions that, on the

one hand, certify the existence of an approximate equilibrium and, on the other, provide guarantees about its efficiency. Then, in subsection 4.2, by use of the Euler–Maclaurin summation formula we present a general form of an approximate potential function, which extends Rosenthal’s potential for weighted congestion games (see also Appendix C). Finally, in subsection 4.3, we deploy this potential for polynomial latencies. Due to its analytic description, our potential differs from other extensions of Rosenthal’s potential that have appeared in previous work, and we believe that this contribution might be of independent interest and could be applied to other classes of latency functions.

**4.1. The potential method.** In the next lemma we lay the foundation for the design and analysis of approximate potential functions by supplying conditions that provide guarantees not only for the existence of approximate equilibria but also for their performance with respect to the social optimum. In the premise of the lemma, we give conditions on the resource functions  $\phi_e$ , having in mind that  $\Phi(\mathbf{s}) = \sum_{e \in E} \phi_e(x_e(\mathbf{s}))$  will eventually serve as the “approximate” potential function.

LEMMA 4.1. *Consider a weighted congestion game with latency functions  $c_e$ , for each facility  $e \in E$ , and player weights  $w_i$ , for each player  $i \in N$ . If there exist functions  $\phi_e : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$  and parameters  $\alpha_1, \alpha_2, \beta_1, \beta_2 > 0$  such that for any facility  $e$  and player weight  $w \in \{w_1, \dots, w_n\}$*

$$(4.1) \quad \alpha_1 \leq \frac{\phi_e(x+w) - \phi_e(x)}{w \cdot c_e(x+w)} \leq \alpha_2 \quad \text{for all } x \geq 0$$

and

$$(4.2) \quad \beta_1 \leq \frac{\phi_e(x)}{x \cdot c_e(x)} \leq \beta_2 \quad \text{for all } x \geq \min_n w_n,$$

then our game has an  $\frac{\alpha_2}{\alpha_1}$ -approximate pure Nash equilibrium which, furthermore, has PoS at most  $\frac{\beta_2}{\beta_1}$ .

*Proof.* Denote  $\alpha = \frac{\alpha_2}{\alpha_1}$ ,  $\beta = \frac{\beta_2}{\beta_1}$ . First we will show that the function  $\Phi(\mathbf{s}) = \sum_{e \in E} \phi_e(x_e(\mathbf{s}))$  (defined over all feasible outcomes  $\mathbf{s}$ ) is an  $\alpha$ -approximate potential; i.e., for any profile  $\mathbf{s}$ , any player  $i$ , and strategy  $s'_i \in S_i$ ,

$$C_i(s'_i, \mathbf{s}_{-i}) < \frac{1}{\alpha} C_i(\mathbf{s}) \implies \Phi(s'_i, \mathbf{s}_{-i}) < \Phi(\mathbf{s}).$$

This would be sufficient to establish the existence of a pure  $\alpha$ -approximate equilibrium, since any (local) minimizer of  $\Phi$  will do. So, it is enough to prove that

$$\Phi(s'_i, \mathbf{s}_{-i}) - \Phi(\mathbf{s}) \leq w_i \alpha_1 [\alpha \cdot C_i(s'_i, \mathbf{s}_{-i}) - C_i(\mathbf{s})].$$

Indeed, if for simplicity we denote  $x_e = x_e(\mathbf{s})$  and  $x'_e = x_e(s'_i, \mathbf{s}_{-i})$  for all facilities  $e$ ,



then we can compute

$$\begin{aligned}
 \Phi(s'_i, \mathbf{s}_{-i}) - \Phi(\mathbf{s}) &= \sum_{e \in E} [\phi_e(x'_e) - \phi_e(x_e)] \\
 &= \sum_{e \in s'_i \setminus s_i} [\phi_e(x_e + w_i) - \phi_e(x_e)] + \sum_{e \in s_i \setminus s'_i} [\phi_e(x_e - w_i) - \phi_e(x_e)] \\
 &\leq \alpha_2 \sum_{e \in s'_i \setminus s_i} w_i c_e(x_e + w_i) - \alpha_1 \sum_{e \in s_i \setminus s'_i} w_i c_e(x_e) \\
 &\leq w_i \alpha_2 \left( \sum_{e \in s'_i \setminus s_i} c_e(x_e + w_i) + \sum_{e \in s'_i \cap s_i} c_e(x_e) \right) \\
 &\quad - w_i \alpha_1 \left( \sum_{e \in s_i \setminus s'_i} c_e(x_e) + \sum_{e \in s'_i \cap s_i} c_e(x_e) \right) \\
 &= w_i \alpha_1 \left[ \alpha \left( \sum_{e \in s'_i \setminus s_i} c_e(x_e + w_i) + \sum_{e \in s'_i \cap s_i} c_e(x_e) \right) \right. \\
 &\quad \left. - \left( \sum_{e \in s_i \setminus s'_i} c_e(x_e) + \sum_{e \in s'_i \cap s_i} c_e(x_e) \right) \right] \\
 &= w_i \alpha_1 [\alpha C_i(s'_i, \mathbf{s}_{-i}) - C_i(\mathbf{s})],
 \end{aligned}$$

where the first inequality holds due to (4.1) and the second holds because  $\alpha_2 \geq \alpha_1$ .

Next, for the upper bound of  $\beta$  on the PoS, it is enough to show that for any profiles  $\mathbf{s}, \mathbf{s}'$ ,

$$\Phi(\mathbf{s}) \leq \Phi(\mathbf{s}') \implies C(\mathbf{s}) \leq \beta \cdot C(\mathbf{s}'),$$

because, if  $\mathbf{s}^* \in \operatorname{argmin}_{\mathbf{s}} C(\mathbf{s})$  is an optimal-cost profile and  $\tilde{\mathbf{s}} \in \operatorname{argmin}_{\mathbf{s}} \Phi(\mathbf{s})$  is a *global* minimizer of  $\Phi$ , then  $C(\tilde{\mathbf{s}}) \leq \beta C(\mathbf{s}^*)$  (and furthermore, as a minimizer of  $\Phi$ ,  $\tilde{\mathbf{s}}$  is clearly an  $\alpha$ -approximate equilibrium as well; see the first part of the current proof). Indeed, denoting  $x_e = x_e(\mathbf{s}), x'_e = x_e(\mathbf{s}')$  for simplicity, we have

$$\begin{aligned}
 \Phi(\mathbf{s}') - \Phi(\mathbf{s}) &= \sum_{e \in E} \phi_e(x'_e) - \sum_{e \in E} \phi_e(x_e) \\
 &\leq \beta_2 \sum_{e \in E} x'_e c_e(x'_e) - \beta_1 \sum_{e \in E} x_e c_e(x_e) \\
 &= \beta_2 C(\mathbf{s}') - \beta_1 C(\mathbf{s}) \\
 &= \beta_1 (\beta C(\mathbf{s}') - C(\mathbf{s})),
 \end{aligned}$$

where for the first inequality we deployed (4.2). □

**4.2. Faulhaber’s potential.** In this section we propose an approximate potential function, which is based on the following classic number-theoretic result, known

as Faulhaber’s formula,<sup>5</sup> which states that for any positive integers  $n, m$ ,

$$\begin{aligned}
 \sum_{k=1}^n k^m &= \frac{1}{m+1} \sum_{j=0}^m (-1)^j \binom{m+1}{j} B_j n^{m+1-j} \\
 (4.3) \qquad &= \frac{1}{m+1} n^{m+1} + \frac{1}{2} n^m + \frac{1}{m+1} \sum_{j=2}^m \binom{m+1}{j} B_j n^{m+1-j},
 \end{aligned}$$

where the coefficients  $B_j$  are the usual Bernoulli numbers.<sup>6</sup> In particular, this shows that the sum of the first  $n$  powers with exponent  $m$  can be expressed as a polynomial of  $n$  with degree  $m+1$ . Furthermore, this sum corresponds to the well-known potential of Rosenthal [46] for *unweighted* congestion games when the latency function is the monomial  $x \mapsto x^m$ .

Based on the above observation, we go beyond just integer values of  $n$  and generalize this idea to all positive reals; in that way, we design a “potential” function that can handle different player weights and, furthermore, incorporate in a more powerful, analytically smooth way approximation factors with respect to both the PoS and the approximation parameter of the equilibrium (in the spirit of Lemma 4.1). A natural way to do this is to directly generalize (4.3) and simply define, for any real  $x \geq 0$  and positive integer  $m$ ,

$$(4.4) \qquad S_m(x) := \frac{1}{m+1} x^{m+1} + \frac{1}{2} x^m,$$

keeping just the first two significant terms.<sup>7</sup> For the special case of  $m = 0$  we set  $S_0(y) := y$ .

For any positive integer  $m$  we define the function  $A_m : [1, \infty) \rightarrow \mathbb{R}_{>0}$  with

$$(4.5) \qquad A_m(x) := \left[ \frac{S_m(x)}{x^{m+1}} \right]^{-1} = \left( \frac{1}{m+1} + \frac{1}{2x} \right)^{-1} = \frac{2(m+1)x}{2x+m+1};$$

for  $m = 0$ , in particular, this gives  $A_0(x) = 1$ . Observe that  $A_m$  is strictly increasing (in  $x$ ) for all  $m \geq 1$ ,

$$(4.6) \qquad A_m(1) = \frac{2(m+1)}{m+3} \in [1, 2), \quad \text{and} \quad \lim_{x \rightarrow \infty} A_m(x) = m+1.$$

For the special case of  $m = 0$  we simply have  $A_0(x) = 1$  for all  $x \geq 0$ . Figure 4.1 shows a graph of these functions. Since  $A_m$  is strictly increasing for  $m \geq 1$ , its inverse function,  $A_m^{-1} : [2\frac{m+1}{m+3}, m+1) \rightarrow [1, \infty)$ , is well defined and also strictly increasing for all  $m \geq 1$ , with

$$(4.7) \qquad A_m^{-1}(x) = \frac{(m+1)x}{2(m+1-x)}.$$

<sup>5</sup>See, e.g., [37, p. 287] or [22, p. 106]). Johann Faulhaber [25] was the first to discover the formula and express it in a systematic way, up to the power of  $m = 17$ . Jakob Bernoulli was able to state it in its full generality as his famous *Summae Potestatum* [7, p. 97] by introducing what are now known as *Bernoulli numbers* (see also footnote 6). The first to rigorously prove the formula was Carl Jacobi [36].

<sup>6</sup>See, e.g., [31, Chapter 6.5] or [1, Chapter 23]. The first Bernoulli numbers are  $B_0 = 1, B_1 = -1/2, B_2 = 1/6, B_3 = 0, B_4 = -1/30, \dots$ . Also, we know that  $B_j = 0$  for all *odd* integers  $j \geq 3$ .

<sup>7</sup>See subsection 4.4 for further discussion on this choice.




FIG. 4.1. Plots of functions  $A_d$  for  $d = 0, 1, 2$  (left) and  $d = 40$  (right). For  $d \geq 1$  they are strictly increasing, starting at  $A_d(1) = \frac{2(d+1)}{d+3} \in [1, 2)$  and going up to  $d + 1$  at the limit. Here,  $A_0(1) = 1$ ,  $A_1(1) = 1$ ,  $A_2(1) = 6/5 = 1.2$ , and  $A_{40}(1) = 82/43 \approx 1.907$ .

The following two lemmas (whose proofs can be found in Appendices B.2 and B.3) describe some useful properties regarding the algebraic behavior of, and the relation among, functions  $A_m$  and  $S_m$ .

LEMMA 4.2. Fix any reals  $y \geq x \geq 1$ . Then the sequences  $\frac{A_m(x)}{m+1}$  and  $\frac{A_m(x)}{A_m(y)}$  are decreasing, and sequence  $A_m(x)$  is increasing (with respect to  $m$ ).

LEMMA 4.3. Fix any integer  $m \geq 0$  and reals  $\gamma, w \geq 1$ . Then

$$(4.8) \quad \frac{\gamma^{m+1}}{A_m(\gamma w)} \leq \frac{S_m(\gamma(x+w)) - S_m(\gamma x)}{w(x+w)^m} \leq \gamma^{m+1} \quad \text{for all } x \geq 0,$$

and

$$(4.9) \quad \frac{\gamma^{m+1}}{m+1} \leq \frac{S_m(\gamma x)}{x^{m+1}} \leq \frac{\gamma^{m+1}}{A_m(\gamma)} \quad \text{for all } x \geq 1.$$

**4.3. The upper bound.** Now we are ready to state our main positive result.

THEOREM 4.4. At any congestion game with polynomial latency functions of degree at most  $d \geq 1$  and player weights ranging in  $[1, W]$ , for any  $\frac{2(d+1)W}{2W+d+1} \leq \alpha \leq d+1$  there exists an  $\alpha$ -approximate pure Nash equilibrium that, furthermore, has PoS at most

$$1 + \left( \frac{d+1}{\alpha} - 1 \right) W.$$

Observe that, as the approximation parameter  $\alpha$  increases, the PoS decreases, in a smooth way from  $\frac{d+3}{2}$  down to the optimal value of 1. Furthermore, notice how the interval within which  $\alpha$  ranges shrinks as the range of player weights  $W$  grows; in particular, its left boundary  $\frac{2(d+1)W}{2W+d+1}$  goes from  $2\frac{d+1}{d+3} = 2 - \frac{4}{d+3}$  (for  $W = 1$ ) up to  $d+1$  (for  $W \rightarrow \infty$ ).

As a result, Theorem 4.4 has two interesting corollaries—one for  $\alpha = \frac{2(d+1)W}{2W+d+1}$  and one for  $W = 1$  (unweighted games).

COROLLARY 4.5. *At any congestion game with polynomial latencies of degree at most  $d \geq 1$  where player weights lie within the range  $[1, W]$ , there is an  $\frac{2(d+1)W}{2W+d+1}$ -approximate pure Nash equilibrium with PoS at most  $\frac{d+3}{2}$ .*

It is interesting to point out here that, in light of Theorem 3.5, the above result of Corollary 4.5 is almost asymptotically tight as far as the PoS is concerned (see the discussion preceding Theorem 3.5).

COROLLARY 4.6. *At any unweighted congestion game with polynomial latencies of degree at most  $d \geq 1$ , the PoS of  $\alpha$ -approximate equilibria is at most  $\frac{d+1}{\alpha}$  for any  $2\frac{d+1}{d+3} \leq \alpha \leq d + 1$ .*

Before proving Theorem 4.4, we first restate it in the following equivalent form that parametrizes the approximation factor of the equilibrium as well as its PoS guarantee, with respect to an “external,” seemingly artificial, parameter  $\gamma \in [1, \infty)$ . The equivalence of the two formulations is formally proven in Appendix B.4.

CLAIM 4.7 (restatement of Theorem 4.4). *For any  $\gamma \geq 1$  there exists an  $A_d(\gamma W)$ -approximate pure Nash equilibrium, which furthermore has PoS at most  $\frac{d+1}{A_d(\gamma)}$ , where  $A_d$  is the strictly increasing function<sup>8</sup> taking values within  $[2\frac{d+1}{d+3}, d + 1)$  defined in (4.5).*

The statement of Claim 4.7 may at first seem a bit cryptic compared to Theorem 4.4. Nevertheless, it brings forth some important aspects of our upper bound construction that are not immediately obvious from Theorem 4.4. In particular, notice how the weight range  $W$  has *no* effect on the PoS guarantee in the statement of Claim 4.7 but appears only in the approximation factor of the equilibrium. Furthermore, as will become more clear in subsection 4.4, this formulation provides a good degree of high-level abstraction that helps with generalizing and improving our result in certain cases, in a unified way. We believe this is important, since it is a promising direction for future work (see also the discussion in Appendix C).

*Proof of Claim 4.7.* Without loss of generality, it is enough to consider only weighted congestion games with *monomial* latency functions (of degree at most  $d$ ); any polynomial is a sum of monomials, so we can just simulate the polynomial latency of a facility by introducing monomial-latency facilities for each one of its summands. More formally, if a facility  $e$  has latency function  $c_e(x) = \sum_{j=0}^d a_{e,j}x^j$ , with constants  $a_{e,0}, a_{e,1}, \dots, a_{e,d} \geq 0$ , we can replace  $e$  by facilities  $e_0, \dots, e_d$  with latencies  $c_{e_j}(x) = a_{e,j}x^j$ , without any change to the costs of the players. Furthermore, we can safely ignore all such facilities  $e_j$  with  $a_{e,j} = 0$ , since they have absolutely no effect on the players’ costs.

So, from now on, assume that for each facility  $e \in E$  there exist a real constant  $a_e > 0$  and a nonnegative integer  $m_e \leq d$  such that

$$c_e(x) = a_e x^{m_e}.$$

Then, in order to utilize Lemma 4.1, we choose functions

$$(4.10) \quad \phi_e(x) = a_e \cdot S_{m_e}(\gamma x),$$

where  $\gamma$  is a real parameter, free to range in  $[1, \infty)$ . Recall here that functions  $S_m$  and  $A_m$  are defined in (4.4) and (4.5). To simplify notation, from now on we fix an arbitrary facility  $e$  and drop the  $e$  subscripts from  $\phi_e, c_e, a_e,$  and  $m_e$ .

<sup>8</sup>See Figure 4.1.

From (4.8) of Lemma 4.3 we get that, for any  $x \geq 0$  and  $w \in [1, W]$ ,

$$\frac{\gamma^{m+1}}{A_m(\gamma w)} \leq \frac{\phi(x+w) - \phi(x)}{w \cdot c(x+w)} = \frac{a [S_m(\gamma(x+w)) - S_m(\gamma x)]}{w \cdot a \cdot (x+w)^m} \leq \gamma^{m+1}.$$

Similarly, from (4.9) we have that for any  $x \geq 1$ ,

$$\frac{\gamma^{m+1}}{m+1} \leq \frac{\phi(x)}{x \cdot c(x)} = \frac{a \cdot S_m(\gamma x)}{x \cdot a \cdot x^m} = \frac{S_m(\gamma x)}{x^{m+1}} \leq \frac{\gamma^{m+1}}{A_m(\gamma)}.$$

Now let us just scale the functions  $\phi_e$  we defined in (4.10) by a factor of  $\frac{1}{S_m(\gamma)}$  and define a potential function

$$\bar{\phi}(x) = \frac{\phi(x)}{S_m(\gamma)} = \frac{A_m(\gamma)}{\gamma^{m+1}} \cdot \phi(x) = a \frac{A_m(\gamma)}{\gamma^{m+1}} S_m(\gamma x).$$

Our previous bounds for  $\phi$  show us that  $\bar{\phi}$  satisfies the requirements of Lemma 4.1 with parameters

$$\begin{aligned} \alpha_1 &= \frac{\gamma^{m+1}}{A_m(\gamma w)} \cdot \frac{A_m(\gamma)}{\gamma^{m+1}} = \frac{A_m(\gamma)}{A_m(\gamma w)} \geq \frac{A_d(\gamma)}{A_d(\gamma w)} \geq \frac{A_d(\gamma)}{A_d(\gamma W)}, \\ \alpha_2 &= \gamma^{m+1} \cdot \frac{A_m(\gamma)}{\gamma^{m+1}} = A_m(\gamma) \leq A_d(\gamma), \\ \beta_1 &= \frac{\gamma^{m+1}}{m+1} \cdot \frac{A_m(\gamma)}{\gamma^{m+1}} = \frac{A_m(\gamma)}{m+1} \geq \frac{A_d(\gamma)}{d+1}, \\ \beta_2 &= \frac{\gamma^{m+1}}{A_m(\gamma)} \cdot \frac{A_m(\gamma)}{\gamma^{m+1}} = 1, \end{aligned}$$

where the inequalities hold due to Lemma 4.2, taking into consideration the fact that  $\gamma w \geq \gamma \geq 1$  and  $m \leq d$ ; specifically, for the last inequality on the bound of  $\alpha_1$ , we also used the fact that  $A_d$  is monotonically increasing.

Putting everything together, from Lemma 4.1 we deduce that indeed there exists an  $A_d(\gamma W)$ -approximate pure Nash equilibrium with PoS at most  $\frac{d+1}{A_d(\gamma)}$ . The fact that  $A_d(\gamma)$  ranges (monotonically) in  $[2^{\frac{d+1}{d+3}}, d+1)$  is a consequence of (4.6).  $\square$

**4.4. Small versus large degree polynomials.** One can argue that our choice to keep only the first two terms in Faulhaber’s formula (4.3), when defining our approximate potential in (4.4), is suboptimal. To some extent, this is correct; it is exactly the reason this seemingly “unnatural” lower bound of  $2^{\frac{d+1}{d+3}} = 2 - \frac{4}{d+3}$  for the approximation parameter  $\alpha$  appears in Corollary 4.6 (or, more generally,  $\frac{2(d+1)W}{2W+d+1}$  in Theorem 4.4). It would be nicer if  $\alpha$  could simply start from 1 instead. Indeed, this can be achieved for small values of  $d$ , as described below.

Considering the entire right-hand side expression in (4.3), one can take the full, exact version of Faulhaber’s formula, which can be written<sup>9</sup> in a very elegant way as

$$(4.11) \quad \sum_{k=1}^n k^m = \frac{1}{m+1} [B_{m+1}(n+1) - B_{m+1}(0)],$$

where

$$B_m(y) = \sum_{k=0}^m \binom{m}{k} B_k y^{m-k}, \quad y \geq 0,$$

<sup>9</sup>See, e.g., [37, p. 288] or [1, eq. 23.1.4].

are the Bernoulli polynomials, and coefficients  $B_k = B_k(0)$  are the standard Bernoulli numbers we used before. Now we can use (4.11) to define a more fine-tuned version for  $S_m$ ; that is, for  $m \geq 1$  we set  $\hat{S}_m(x) = \frac{1}{m+1} [B_{m+1}(x+1) - B_{m+1}]$  instead of (4.4). For example, for degrees up to  $m \leq 4$  these new polynomials are

$$\begin{aligned} \hat{S}_0(x) &= x, & \hat{S}_1(x) &= \frac{1}{2}x(x+1), & \hat{S}_2(x) &= \frac{1}{6}x(2x^2 + 3x + 1), \\ \hat{S}_3(x) &= \frac{1}{4}x^2(x+1)^2, & \hat{S}_4(x) &= \frac{1}{30}x(6x^4 + 15x^3 + 10x^2 - 1). \end{aligned}$$

Using these values, one can verify that for up to  $m \leq 4$ , all our critical technical requirements for the proof of Claim 4.7 (and thus, Theorem 4.4 itself) are satisfied, most notably Lemmas 4.2 and 4.3 and the monotonicity of  $\hat{A}_m(x) = \frac{x^{m+1}}{\hat{S}_m(x)}$  (with respect to  $x \geq 1$ ). In particular, now we have that  $\hat{A}_m(1) = \frac{1^{m+1}}{\hat{S}_m(1)} = 1$ , which is exactly what we wanted: this means that the critical quantities  $\hat{A}_d(\gamma W)$  and  $\hat{A}_d(\gamma)$  in Claim 4.7 can start taking values all the way down to  $\hat{A}_d(W)$  and  $\hat{A}_d(1) = 1$ , respectively. This translates to the approximation ratio parameter  $\alpha$  in our main result in Theorem 4.4 starting to range from  $\alpha \geq \hat{A}_d(W)$ .

Thus,

*Theorem 4.4 can be rewritten for  $d \leq 4$ , with the approximation parameter  $\alpha$  taking values in  $\hat{A}_d(W) \leq \alpha \leq d + 1$ . In particular, for unweighted games, this means that Corollary 4.6 can be rewritten with  $\alpha$  taking values within the entire range of  $[1, d + 1]$ .*

However, there is a catch which does not allow us to do that in general; as  $m$  grows large, the Bernoulli polynomials, which now play a critical role in our definition of functions  $\hat{S}_m$  (see (4.11)), start to behave in a rather erratic, nonsmooth way within the interior of the real intervals between consecutive integer values. For example, one can check that, for  $d = 14$ , function  $\hat{A}_{14}$  is *not* monotonically increasing within  $[1, 2]$ . Even more disastrously, for  $d = 20, 21$ , functions  $\hat{S}_d$  take *negative* values in  $[1, 2]$ !

**Appendix A. Lower bound proofs.**

**A.1. Technical lemmas.**

LEMMA A.1. *For any  $d \geq 9$ ,*

$$\left(1 + \frac{\ln d}{d}\right)^d \geq \frac{d}{\ln d}.$$

*Proof.* From Mitrinović [42, eq. (3), p. 267] we know that the following inequality holds for all  $n \geq 1$  and  $1 \leq x \leq n$ :

$$\left(1 + \frac{x}{n}\right)^n \geq e^x \left(1 - \frac{x^2}{n}\right).$$

Applying it with  $n = d$  and  $x = \ln d$ , we get that indeed

$$\left(1 + \frac{\ln d}{d}\right)^d \geq e^{\ln d} \left(1 - \frac{\ln^2 d}{d}\right) \geq d \frac{1}{\ln d},$$

the last inequality holding due to the fact that for  $d \geq 9$ ,  $\frac{\ln^2 d}{d} + \frac{1}{\ln d} \leq \frac{\ln^2(9)}{9} + \frac{1}{\ln(9)} \approx 0.992 \leq 1$ . □

LEMMA A.2. For any integer  $d \geq 2$ , the function  $f : (0, \infty)^2 \rightarrow (0, \infty)$  defined by

$$f(x, y) = \frac{(y + x + 1)^d - (y + x)^d}{(y + 1)^d - y^d}$$

is monotonically decreasing with respect to  $y$ . Furthermore, for  $d \geq 9$ ,

$$(A.1) \quad \zeta^{d+1} \leq f((\beta_d \Phi_d + 1)(\zeta - 1), \beta_d \Phi_d - (1 - \beta_d)\zeta) \quad \text{for all } \zeta \in [1, 2],$$

where  $\beta_d$  is defined in Lemma 3.2.

*Proof.* First, let us define function  $h : (0, \infty) \rightarrow (0, \infty)$  with

$$(A.2) \quad h(t) = \frac{(t + 1)^d - t^d}{(t + 1)^{d-1} - t^{d-1}}.$$

We will show that  $h$  is increasing, which will suffice to prove the desired monotonicity of  $f$  since its derivative is

$$\begin{aligned} \frac{\partial f(x, y)}{\partial y} &= \frac{d [(x + y + 1)^{d-1} - (x + y)^{d-1}]}{(y + 1)^d - y^d} \\ &\quad - \frac{d [(y + 1)^{d-1} - y^{d-1}] [(x + y + 1)^d - (x + y)^d]}{[(y + 1)^d - y^d]^2} \\ &= \frac{d [(y + 1)^{d-1} - y^{d-1}] [(x + y + 1)^{d-1} - (x + y)^{d-1}]}{[(y + 1)^d - y^d]^2} [h(y) - h(x + y)], \end{aligned}$$

which is negative due to the monotonicity of  $h$ . To prove that  $h$  is indeed increasing, we will show something stronger—namely that function  $\bar{h} : (1, \infty) \rightarrow (0, \infty)$  with

$$(A.3) \quad \bar{h}(t) = \frac{t^d - (t - 1)^d}{t^d - t(t - 1)^{d-1}}$$

is increasing. This will suffice to demonstrate that  $h$  is increasing as well, since  $h(t) = (t + 1) \cdot \bar{h}(t + 1)$ . Taking its derivative, we see that

$$\frac{\partial \bar{h}(t)}{\partial t} = \frac{[(t - 1)^d - t^d + dt^{d-1}] (t - 1)^d}{[t^{d+1} - t^d - t(t - 1)^d]^2} > 0$$

since from the convexity of function  $t \mapsto t^d$  we know that  $t^d - (t - 1)^d < dt^{d-1}$ .

Now let us prove the remaining part of our lemma, that is, (A.1). Observe that if we set  $\zeta = 1$  to (A.1), it is satisfied, since  $f(0, y) = 1$  for any  $y > 0$ . So, it is enough if we prove that

$$\begin{aligned} &\zeta^{-(d+1)} f((\beta_d \Phi_d + 1)(\zeta - 1), \beta_d \Phi_d - (1 - \beta_d)\zeta) \\ &= \zeta^{-(d+1)} \frac{[(\alpha + \beta)\zeta]^d - [(\alpha + \beta)\zeta - 1]^d}{[\alpha + 1 - (1 - \beta)\zeta]^d - [\alpha - (1 - \beta)\zeta]^d} \end{aligned}$$

is increasing with respect to  $\zeta \in [1, 2]$ , where here we are using  $\beta = \beta_d$  and  $\alpha = \beta \Phi_d$ . So, if we define

$$\begin{aligned} f_1(\zeta) &= [(\alpha + \beta)\zeta]^d - [(\alpha + \beta)\zeta - 1]^d, \\ f_2(\zeta) &= [\alpha + 1 - (1 - \beta)\zeta]^d - [\alpha - (1 - \beta)\zeta]^d \end{aligned}$$

and we compute the derivative  $\frac{\partial}{\partial \zeta}(\zeta^{-(d+1)} \frac{f_1(\zeta)}{f_2(\zeta)})$  of the above expression, we need to show that

$$(A.4) \quad \zeta \left[ \frac{f'_1(\zeta)}{f_1(\zeta)} - \frac{f'_2(\zeta)}{f_2(\zeta)} \right] \geq d + 1.$$

Now notice that

$$\zeta \frac{f'_1(\zeta)}{f_1(\zeta)} = d(\alpha + \beta)\zeta \frac{[(\alpha + \beta)\zeta]^{d-1} - [(\alpha + \beta)\zeta - 1]^{d-1}}{[(\alpha + \beta)\zeta]^d - [(\alpha + \beta)\zeta - 1]^d} = \frac{d}{\bar{h}((\alpha + \beta)\zeta)},$$

where  $\bar{h}$  is the increasing function defined in (A.3), so taking into consideration that

$$(\alpha + \beta)\zeta = \beta(\Phi_d + 1)\zeta \leq \frac{1}{2}(\Phi_d + 1)2 \leq \Phi_d + 1,$$

we can get that

$$\begin{aligned} \zeta \frac{f'_1(\zeta)}{f_1(\zeta)} &\geq \frac{d}{\bar{h}(\Phi_d + 1)} = d(\Phi_d + 1) \frac{(\Phi_d + 1)^{d-1} - \Phi_d^{d-1}}{(\Phi_d + 1)^d - \Phi_d^d} \\ &= d \frac{\Phi_d^{d+1} - \Phi_d^d - \Phi_d^{d-1}}{\Phi_d^{d+1} - \Phi_d^d} = d - \frac{d}{\Phi_d^2 - \Phi_d}. \end{aligned}$$

Similarly, we can see that

$$-\zeta \frac{f'_2(\zeta)}{f_2(\zeta)} = \frac{d(1 - \beta)\zeta}{h(\alpha - (1 - \beta)\zeta)},$$

where  $h$  is the increasing function defined in (A.2), so taking into consideration that

$$\alpha - (1 - \beta)\zeta \leq \beta\Phi_d - (1 - \beta) \leq \frac{\Phi_d - 1}{2} \quad \text{and} \quad (1 - \beta)\zeta \geq \frac{1}{2},$$

we get that

$$-\zeta \frac{f'_2(\zeta)}{f_2(\zeta)} \geq \frac{d/2}{h((\Phi_d - 1)/2)} = d \frac{(\Phi_d + 1)^{d-1} - (\Phi_d - 1)^{d-1}}{(\Phi_d + 1)^d - (\Phi_d - 1)^d}.$$

Putting everything together, in order to prove the desired (A.4), it now suffices to show that

$$d \frac{(\Phi_d + 1)^{d-1} - (\Phi_d - 1)^{d-1}}{(\Phi_d + 1)^d - (\Phi_d - 1)^d} - \frac{d}{\Phi_d^2 - \Phi_d} \geq 1,$$

which we know holds from (A.5) of Lemma A.3. □

LEMMA A.3. *For any integer  $d \geq 9$ ,*

$$(A.5) \quad \frac{(\Phi_d + 1)^{d-1} - (\Phi_d - 1)^{d-1}}{(\Phi_d + 1)^d - (\Phi_d - 1)^d} - \frac{1}{\Phi_d^2 - \Phi_d} \geq \frac{1}{d}.$$

Furthermore, asymptotically  $\Phi_d \sim \frac{d}{\ln d}$ , i.e.,

$$(A.6) \quad \lim_{d \rightarrow \infty} \frac{\Phi_d}{d/\ln d} = 1.$$



In particular, for any integer  $d$ ,

$$(A.7) \quad \Phi_d \leq \gamma_d \frac{d}{\ln d} \quad \text{with} \quad \gamma_d := \frac{\ln d}{\mathcal{W}(d)} \leq 1.368 \quad \text{and} \quad \lim_{d \rightarrow \infty} \gamma_d = 1,$$

where  $\mathcal{W}(\cdot)$  denotes the (principal branch of the) Lambert- $W$  function.<sup>10</sup>

*Proof.* To show (A.5), first one can simply numerically verify that it indeed holds for all integers  $d = 9, 10, \dots, 17$ , so let us just focus on the case when  $d \geq 18$ . For simplicity, in the remainder of the proof we denote  $y = \Phi_d$ . It is easy to see<sup>11</sup> then that

$$y \geq \Phi_{18} \approx 8.11 > 8.$$

Performing some elementary algebraic manipulations in (A.5), we can equivalently write it as

$$(y + 1)^{d-1} [-y^3 + dy^2 - (2d - 1)y - d] \geq (y - 1)^{d+1}(d - y).$$

Using the fact that  $(y + 1)^{d-1} = \frac{y^{d+1}}{y+1}$ , and then that  $y^{d+1} \geq (y - 1)^{d+1}$ , we can see that it is enough to show that

$$-y^3 + dy^2 - (2d - 1)y - d \geq (d - y)(y + 1)$$

or, equivalently,

$$(A.8) \quad \begin{aligned} y^2(d - y) - 2dy + y - d &\geq (d - y)(y + 1), \\ (d - y)[y^2 - 1 - (y + 1)] &\geq 2dy, \\ (d - y)(y + 1)(y - 2) &\geq 2yd, \\ \frac{(y + 1)(y - 2)}{2y} &\geq \frac{1}{1 - \frac{y}{d}}. \end{aligned}$$

For the last inequality we took into consideration that  $\frac{y}{d} < 1$ . As a matter of fact, using an upper bound of  $y \leq \frac{2d}{\ln d}$  on  $y = \Phi_d$  (see [3, Lemma 5.4] or (A.7)), we have that

$$\frac{y}{d} \leq \frac{2}{\ln d} \leq \frac{2}{\ln(18)} \approx 0.692.$$

Due to the fact that function  $z \mapsto \frac{1}{1-z}$  is increasing for  $z \in [0, 1)$ , this bound gives us that  $\frac{1}{1-\frac{y}{d}} \leq \frac{1}{1-0.69} \leq 3.226$ . In a similar way, noticing that function  $z \mapsto \frac{(z+1)(z-2)}{2z}$  is increasing for  $z > 2$ , using the fact that  $y > 8$ , we can derive that  $\frac{(y+1)(y-2)}{2y} > \frac{(8+1)(8-2)}{2 \cdot 8} = \frac{27}{8} \approx 3.375$ . This establishes the validity of (A.8).

Next, we deal with upper-bounding the values of  $\Phi_d$  and proving (A.7), since we will need this for establishing the asymptotics of  $\Phi_d$  in (A.6). Here we will make use of the following property, which was shown in Aland et al. [3, Lemma 5.2, Theorem 3.4]: for any real  $\gamma > 0$ ,

$$(A.9) \quad \Phi_d < \gamma \frac{d}{\ln d} \iff \left(1 + \frac{\ln d}{\gamma d}\right)^d < \frac{\gamma d}{\ln d}.$$

<sup>10</sup>That is, for any positive real  $x$ ,  $\mathcal{W}(x) = z$  gives the unique positive real solution  $z$  to the equation  $x = z \cdot e^z$ .

<sup>11</sup>Here we are silently using the fact that  $\Phi_n$  is an increasing function of the integer  $n$ . One can formally prove this by, e.g., combining Lemmas 5.1 and 5.2 of Aland et al. [3].

In a similar way to the proof of Aland et al. [3, Theorem 3.4], using a binomial expansion, we can compute

$$\begin{aligned} \left(1 + \frac{\ln d}{\gamma d}\right)^d &= \sum_{k=0}^d \binom{d}{k} \left(\frac{\ln d}{\gamma d}\right)^k = \sum_{k=0}^d \frac{d!}{d^k (d-k)!} \left(\frac{\ln d}{\gamma}\right)^k \frac{1}{k!} \\ &= \sum_{k=0}^d \left(1 - \frac{1}{d}\right) \left(1 - \frac{2}{d}\right) \dots \left(1 - \frac{k-1}{d}\right) \left(\frac{\ln d}{\gamma}\right)^k \frac{1}{k!} \\ &< \sum_{k=0}^{\infty} \frac{\left(\frac{\ln d}{\gamma}\right)^k}{k!} = e^{\frac{\ln d}{\gamma}} = d^{1/\gamma}, \end{aligned}$$

where for the second to last equality we used the power series representation of the exponential function:  $e^z = \sum_{k=0}^{\infty} \frac{z^k}{k!}$ . Thus, from (A.9) we can derive that, for any  $\gamma > 0$ ,

$$(A.10) \quad d^{1/\gamma} \leq \frac{\gamma \cdot d}{\ln d} \implies \Phi_d < \gamma \frac{d}{\ln d}.$$

Using  $\gamma_d = \frac{\ln d}{\mathcal{W}(d)}$  as defined in the statement of our lemma, we compute

$$d^{1/\gamma_d} = d^{\frac{\mathcal{W}(d)}{\ln d}} = e^{\mathcal{W}(d)}$$

and

$$\gamma_d \frac{d}{\ln d} = \frac{\ln d}{\mathcal{W}(d)} \frac{d}{\ln d} = \frac{d}{\mathcal{W}(d)}.$$

Thus, since from the definition of function  $\mathcal{W}$  we know that

$$(A.11) \quad \mathcal{W}(d)e^{\mathcal{W}(d)} = d,$$

we deduce that  $\gamma = \gamma_d$  indeed satisfies the left-hand side of (A.10), giving us the desired upper bound for  $\Phi_d$ .

For the asymptotic behavior of  $\gamma_d$  when  $d$  grows large, observe that by taking logarithms in (A.11) we get

$$\mathcal{W}(d) + \ln \mathcal{W}(d) = \ln d,$$

and so

$$\lim_{d \rightarrow \infty} \gamma_d = \lim_{d \rightarrow \infty} \frac{\ln d}{\mathcal{W}(d)} = \lim_{d \rightarrow \infty} \left[ \frac{\ln \mathcal{W}(d)}{\mathcal{W}(d)} + 1 \right] = 1 + \lim_{z \rightarrow \infty} \frac{\ln z}{z} = 1,$$

since it is easy to see that  $\lim_{d \rightarrow \infty} \mathcal{W}(d) = \infty$ .

Finally, let us now establish (A.6). Due to (A.7), which we have already proved, it is enough to just show a lower bound of  $\lim_{d \rightarrow \infty} \frac{\Phi_d}{d/\ln d} \geq 1$ . We will do this by showing that  $\Phi_d \geq \frac{d}{\ln d}$  for sufficiently large values of  $d$ . Indeed, by (A.9) this is equivalent to proving that

$$\left(1 + \frac{\ln d}{d}\right)^d \geq \frac{d}{\ln d},$$

which from Lemma A.1 we know holds for all  $d \geq 9$ . □

**A.2. Proof of Lemma 3.2.** To decongest notation a bit in the proof, we will drop the  $d$  subscripts from  $c_d$  and  $\beta_d$  whenever this causes no confusion. Starting with (3.3), if we solve with respect to  $\beta$ , we get

$$(A.12) \quad \Phi_d + \frac{1}{\beta} \geq \Phi_d^{1+\frac{2}{d}} \iff \beta \leq \frac{1}{\Phi_d(\Phi_d^{2/d} - 1)} = \frac{\Phi_d}{2\Phi_d + 1},$$

where for the last equality we used the fact that

$$\Phi_d^{2/d} = \left(1 + \frac{1}{\Phi_d}\right)^2 = \frac{1}{\Phi_d^2} + \frac{2}{\Phi_d} + 1,$$

which is a direct consequence of the definition of  $\Phi_d$ :

$$\Phi_d^{d+1} = (\Phi_d + 1)^d \iff \Phi_d^{1+\frac{1}{d}} = \Phi_d + 1 \iff \Phi_d^{1/d} = 1 + \frac{1}{\Phi_d}.$$

Substituting (3.2) into (A.12), we have

$$\begin{aligned} 1 - \Phi_d^{-c} \leq \frac{\Phi_d}{2\Phi_d + 1} &\iff \Phi_d^{-c} \geq \frac{\Phi_d + 1}{2\Phi_d + 1} \\ &\iff c \leq \frac{\ln(2\Phi_d + 1) - \ln(\Phi_d + 1)}{\ln \Phi_d}, \end{aligned}$$

which holds by the very definition of  $c$  in (3.1) if we relax the floor operator.

Let us now move to (3.4) and in particular lower-bound the values of parameter  $\beta$  as  $d$  grows large. Due to the floor operator in (3.1), parameter  $c$  can be lower-bounded by

$$c \geq \frac{\ln(2\Phi_d + 1) - \ln(\Phi_d + 1)}{\ln \Phi_d} - \frac{1}{d} = \log_{\Phi_d} \frac{2\Phi_d + 1}{\Phi_d + 1} - \frac{1}{d},$$

and since  $\beta$  is increasing with respect to  $c$ ,

$$(A.13) \quad \beta = 1 - \Phi_d^{-c} \geq 1 - \frac{\Phi_d + 1}{2\Phi_d + 1} \Phi_d^{1/d} = 1 - \frac{\Phi_d + 1}{2\Phi_d + 1} \left(1 + \frac{1}{\Phi_d}\right).$$

Taking limits and recalling that  $\lim_{d \rightarrow \infty} \Phi_d = \infty$ , we get the desired lower bound of

$$\lim_{d \rightarrow \infty} \beta \geq 1 - \frac{1}{2} \cdot (1 + 0) = \frac{1}{2}.$$

The upper bound of  $\beta \leq \frac{1}{2}$  can be easily derived by (A.12):

$$\beta \leq \frac{\Phi_d}{2\Phi_d + 1} \leq \frac{\Phi_d}{2\Phi_d} = \frac{1}{2}.$$

For small values of  $d$ , and in particular in order to prove that  $\beta \geq 0.38$ , one can numerically compute the values for  $\beta$  directly from (3.1). For example, for  $d = 9, \dots, 100$ , these values are shown in Figure 3.1. The lower and upper red lines in Figure 3.1 correspond to the relaxation of the floor operator we used in the lower and upper bounds for  $\beta$  in (A.13) and (A.12), respectively. The actual values of  $\beta$  lie between these two lines. Using these values and the resulting monotonicity for  $\beta$ , one can also prove the lower bound of 3 for  $dc$  by observing that, by setting  $d = 9$  in (3.1), we have that for any  $d \geq 9$

$$d \cdot c \geq \left\lceil 9 \frac{\ln(2 \cdot \Phi_9 + 1) - \ln(\Phi_9 + 1)}{\ln(\Phi_9)} \right\rceil \approx \lceil 3.368 \rceil = 3,$$

since  $\Phi_9 \approx 5.064$ .

**A.3. Proof of Claim 3.3.** Since we have fixed that  $s_j = \tilde{s}_j$  for all players  $j < i$  and also the strategies of players  $j > i + \mu$  have no effect on the cost of player  $i$  (and in particular in (3)), it is safe to briefly abuse notation and for now assume that  $\mathbf{s}_{-i} = (s_{i+1}, \dots, s_{i+\mu})$ .

We generate the desired dominating profile  $\mathbf{s}'_{-i}$  inductively by running Procedure  $\text{DOMINATE}(\mathbf{s}_{-i}, i)$ , described formally below, scanning and modifying profile  $\mathbf{s}_{-i}$  from right to left.

---

**Procedure**  $\text{DOMINATE}(\mathbf{s}_{-i}, i)$

---

**Input:** Profile  $\mathbf{s}_{-i} = (s_{i+1}, \dots, s_{i+\mu})$ ; Player  $i \in \{\mu + 1, \dots, n\}$   
**Output:** Profile  $\mathbf{s}'_{-i} = (s'_{i+1}, \dots, s'_{i+\mu})$  of the form described in Items 1 to 3 of Claim 3.3, which satisfies (3.7)

- 1  $\mathbf{s}'_{-i} \leftarrow \mathbf{s}_{-i}$ ;
- 2  $s'_{i+\mu} \leftarrow s^*_{i+\mu}$ ;
- 3  $k \leftarrow i + \mu - 1$ ;
- 4 **while** exists  $j \in \{i + 1, \dots, k - 1\}$  such that  $s'_j = s^*_j$  **do**
- 5      $s'_k \leftarrow \tilde{s}_k$ ;
- 6      $k \leftarrow k - 1$ ;

**end**

---

First, it is not difficult to see that the output profile  $\mathbf{s}'_{-i}$  of  $\text{DOMINATE}(\mathbf{s}_{-i}, i)$  indeed has the desired format described in Items 1 to 3 of Claim 3.3. In particular, after any execution of the while-loop in lines 4–6 of Procedure  $\text{DOMINATE}$ ,  $s'_j = \tilde{s}_j$  for any  $j = k + 1, \dots, i + \mu - 1$ . Furthermore, it is also easy to see that switching player’s  $i + \mu$  strategy to  $s'_{i+\mu} = s^*_{i+\mu}$  can only increase player’s  $i$  cost, i.e., (3.7) is satisfied after line 2 of  $\text{DOMINATE}$ : if player  $i + \mu$  chooses  $\tilde{s}_{i+\mu}$  instead, she contributes nothing to the cost of player  $i$ , since she does not put her weight in any of the facilities  $i + 1, \dots, i + \mu$  played by player  $i$ .

So, it remains to be shown that after every iteration of the while-loop, condition (3.7) is maintained. Since in any such loop only the strategy of player  $k$  is possibly switched from  $s^*_k$  to  $\tilde{s}_k$ , it is enough if we show that  $C_i(\tilde{s}_k, \mathbf{s}'_{-k}) \geq C_i(s^*_k, \mathbf{s}'_{-k})$  or, since for any facility  $j < k$  it holds that  $x_j(\tilde{s}_k, \mathbf{s}'_{-k}) = x_j(s^*_k, \mathbf{s}'_{-k})$ , equivalently

$$\sum_{j=k}^{i+\mu} c_j(x_j(\tilde{s}_k, \mathbf{s}'_{-k})) \geq \sum_{j=k}^{i+\mu} c_j(x_j(s^*_k, \mathbf{s}'_{-k})).$$

If we let  $z_j$ , for any  $j \geq k$ , denote the load on facility  $j$  induced by every player *except* from player  $k$ , that is, formally,

$$z_j = \sum \{w_\ell \mid \ell \in \{j - \mu, \dots, j\} \setminus \{k\} \wedge j \in s'_\ell\},$$

then the above can be written as

$$\sum_{j=k+1}^{i+\mu} [c_j(z_j + w_k) - c_j(z_j)] \geq c_k(z_k + w_k) - c_k(z_k).$$

Thus, it is sufficient to only take  $j = i + \mu$  in the above sum and just prove that

$$c_{i+\mu}(z_{i+\mu} + w_k) - c_{i+\mu}(z_{i+\mu}) \geq c_k(z_k + w_k) - c_k(z_k),$$

which is equivalent to

$$(A.14) \quad w^{-(i+\mu-k)(d+1)} \frac{(z_{i+\mu} + w_k)^d - z_{i+\mu}^d}{(z_k + w_k)^d - z_k^d} \geq 1.$$

If we define

$$A = \{j \in \{i + 1, \dots, k - 1\} \mid s_j = s_j^*\},$$

that is,  $A$  is the set of players below  $k$  (and above  $i$ ) that do *not* contribute with their weight to the cost of players  $k$  and  $i + \mu$ , then we have that

$$z_k = \sum_{\substack{j=k-\mu \\ j \notin A}}^{k-1} w_j = \alpha w_k - \sum_{j \in A} w_j$$

and

$$z_{i+\mu} = \sum_{\substack{j=i \\ j \in A \cup \{k\}}}^{i+\mu} w_j = (\alpha + 1)w_{i+\mu} - w_k - \sum_{j \in A} w_j,$$

because by our inductive process we know that  $s'_j = \tilde{s}_j$  for all  $j = k + 1, \dots, i + \mu - 1$ . Thus

$$z_{i+\mu} = z_k + (\alpha + 1)(w_{i+\mu} - w_k).$$

Now we can rewrite the left-hand side of (A.14) as

$$w^{-(i+\mu-k)(d+1)} \frac{[z_k + (\alpha + 1)w_{i+\mu} - \alpha w_k]^d - [z_k + (\alpha + 1)w_{i+\mu} - (\alpha + 1)w_k]^d}{(z_k + w_k)^d - z_k^d},$$

and, if we additionally define for simplicity

$$\zeta := w^\lambda, \quad \text{where} \quad \lambda := \mu - k + i \in \{1, \dots, \mu - 2\}$$

and

$$y_k := \frac{z_k}{w_k},$$

then (A.14) can be written as

$$\zeta^{-(d+1)} \frac{[y_k + (\alpha + 1)\zeta - \alpha]^d - [y_k + (\alpha + 1)\zeta - \alpha - 1]^d}{(y_k + 1)^d - y_k^d} \geq 1,$$

or, more simply,

$$\zeta^{-(d+1)} f(x, y) \geq 1$$

if we use function  $f$  from Lemma A.2 with values

$$x = (\alpha + 1)w^\lambda - \alpha - 1 = (\alpha + 1)(\zeta - 1) > 0 \quad \text{and} \quad y = y_k > 0.$$

Deploying the monotonicity of  $f$  from Lemma A.2 and using that

$$y = \frac{z_k}{w_k} = \alpha - \frac{1}{w_k} \sum_{j \in A} w_j \leq \alpha - \frac{1}{w_k} w_{i+1} = \alpha - w^{i+1-k} = \alpha - w^{\lambda-\mu+1} \leq \alpha - (1 - \beta)\zeta,$$

where the first inequality holds due to the fact that from the while-loop test in line 4 of Procedure DOMINATE we know that  $A \neq \emptyset$  and the last one holds because of  $1 - \beta = w^{-\mu}$ , we finally get that it is enough if we show that

$$\zeta^{d+1} \leq f((\alpha + 1)(\zeta - 1), \alpha - (1 - \beta)\zeta) \quad \text{for all } \zeta \in \left[1, \frac{1}{1 - \beta}\right],$$

since  $\zeta \geq w^1 \geq 1$  and  $\zeta \leq w^{\mu-2} \leq w^\mu = (1 - \beta)^{-1}$ . But this is satisfied, due to (A.1) of Lemma A.2, since we have selected our parameters  $c = c_d$  and  $\beta = \beta_d$  as in Lemma 3.2.

**A.4. Proof of (3.8).** In any such profile, player  $i + \mu$  plays  $s_{i+\mu}^*$  and the following hold:

- Either all other players  $j = i + 1, \dots, i + \mu - 1$  play  $\tilde{s}_j$ , in which case

$$\begin{aligned} C_i(\tilde{s}_i, \mathbf{s}'_{-i}) &= c_{i+\mu} \left( \sum_{\ell=i}^{i+\mu} w_\ell \right) + \sum_{j=i+1}^{i+\mu-1} c_j \left( \sum_{\ell=j-\mu}^{j-1} w_\ell \right) \\ &= c_{i+\mu} ((\alpha + 1)w^{i+\mu}) + \sum_{j=i+1}^{i+\mu-1} c_j (\alpha w^j) \\ &= w^{-(i+\mu)(d+1)} (\alpha + 1)^d w^{d(i+\mu)} + \sum_{j=i+1}^{i+\mu-1} w^{-j(d+1)} \alpha^d w^{dj} \\ &= w^{-(i+\mu)} (\alpha + 1)^d + \sum_{j=i+1}^{i+\mu-1} w^{-j} \alpha^d \\ &= w^{-i} \left[ (\alpha + 1)^d w^{-\mu} + \alpha^d \sum_{j=1}^{\mu-1} w^{-j} \right] \\ &= w^{-i} [(\alpha + 1)^d w^{-\mu} + \alpha^d (\alpha - w^{-\mu})], \end{aligned}$$

the last equality holding due to the definition of  $\alpha$ ,

- or there exists a *single* player  $k \in \{i + 1, \dots, i + \mu - 1\}$  that plays  $s_k^*$  (instead of  $\tilde{s}_k$ , which corresponds exactly to the previous case), in which case

$$\begin{aligned} C_i(\tilde{s}_i, \mathbf{s}'_{-i}) &\leq c_k \left( \sum_{\ell=k-\mu}^k w_\ell \right) + c_{i+\mu} \left( \sum_{\ell=i}^{i+\mu} w_\ell - w_k \right) + \sum_{\substack{j=i+1 \\ j \neq k, i+\mu}}^{i+\mu} c_j \left( \sum_{\ell=j-\mu}^{j-1} w_\ell \right) \\ &= c_k ((\alpha + 1)w^k) + c_{i+\mu} ((\alpha + 1)w^{i+\mu} - w^k) + \sum_{\substack{j=i+1 \\ j \neq k}}^{i+\mu-1} c_j (\alpha w^j) \\ &= w^{-i} \left[ (\alpha + 1)^d w^{-(k-i)} + (\alpha + 1 - w^{k-i-\mu})^d w^{-\mu} \right. \\ &\quad \left. + \alpha^d (\alpha - w^{-\mu} - w^{-(k-i)}) \right], \end{aligned}$$

which is decreasing with respect to  $k$ , so taking the smallest possible value  $k = i + 1$  we have that

$$\begin{aligned} C_i(\tilde{s}_i, \mathbf{s}'_{-i}) &\leq w^{-i} [(\alpha + 1)^d w^{-1} + (\alpha + 1 - w^{1-\mu})^d w^{-\mu} + \alpha^d (\alpha - w^{-\mu} - w^{-1})]. \end{aligned}$$

Considering both of the above possible scenarios, in order to prove (3.8) it is thus sufficient to make sure that

$$(A.15) \quad \alpha^d(\alpha - w^{-\mu}) < (\alpha + 1)^d(1 - w^{-\mu})$$

and

$$(A.16) \quad (\alpha + 1 - w^{1-\mu})^d w^{-\mu} + \alpha^d(\alpha - w^{-\mu} - w^{-1}) < (\alpha + 1)^d(1 - w^{-1}).$$

For (A.15), its left-hand side can be written as

$$(\beta\Phi_d)^d (\beta\Phi_d - (1 - \beta)) < (\beta\Phi_d)^d (\beta\Phi_d) = \beta^{d+1}\Phi_d^{d+1} = \beta^{d+1}(\Phi_d + 1)^d,$$

the first inequality holding because  $\beta < 1$ , while the right-hand side is

$$(\beta\Phi_d + 1)^d(1 - (1 - \beta)) = \beta^{d+1} \left( \Phi_d + \frac{1}{\beta} \right)^d.$$

Thus it is enough to prove that

$$(\Phi_d + 1)^d \leq \left( \Phi_d + \frac{1}{\beta} \right)^d,$$

which holds since  $\beta \in (0, 1)$ .

For (A.16), the left-hand side is written as

$$\begin{aligned} & \left( \beta\Phi_d + 1 - \left( 1 + \frac{1}{\Phi_d} \right) (1 - \beta) \right)^d (1 - \beta) \\ & \quad + (\beta\Phi_d)^d \left( \beta\Phi_d - (1 - \beta) - \left( 1 + \frac{1}{\Phi_d} \right)^{-1} \right) \\ & = \left( \beta\Phi_d + \beta - \frac{1 - \beta}{\Phi_d} \right)^d (1 - \beta) + \beta^d \Phi_d^d \left( \beta\Phi_d - (1 - \beta) - \frac{\Phi_d}{\Phi_d + 1} \right) \\ & < (\beta\Phi_d + \beta)^d (1 - \beta) + \beta^d \Phi_d^d \left( \beta\Phi_d - \frac{\Phi_d}{\Phi_d + 1} \right) \\ & = \beta^d (\Phi_d + 1)^d (1 - \beta) + \beta^d \Phi_d^d \left( \beta\Phi_d - \frac{\Phi_d}{\Phi_d + 1} \right) \\ & = \beta^d (\Phi_d)^{d+1} (1 - \beta) + \beta^d \Phi_d^d \left( \beta\Phi_d - \frac{\Phi_d}{\Phi_d + 1} \right) \\ & = \beta^d \Phi_d^d \left( \Phi_d - \beta\Phi_d + \beta\Phi_d - \frac{\Phi_d}{\Phi_d + 1} \right) \\ & = \beta^d \frac{\Phi_d^{d+2}}{\Phi_d + 1}, \end{aligned}$$

and the right-hand side is written as

$$(\beta\Phi_d + 1)^d \left( 1 - \left( 1 + \frac{1}{\Phi_d} \right)^{-1} \right) = \beta^d \left( \Phi_d + \frac{1}{\beta} \right)^d \frac{1}{\Phi_d + 1}.$$

Thus it suffices to prove that

$$\Phi_d^{d+2} \leq \left( \Phi_d + \frac{1}{\beta} \right)^d,$$

which holds due to (3.3) since we have already selected parameter  $c = c_d$  as in (3.1).

**Appendix B. Upper bound proofs.**

**B.1. Technical lemmas.**

LEMMA B.1. *For any positive integer  $m$  and real  $x > 0$ ,*

$$\left(1 + \frac{1}{x}\right)^m \geq 1 + \frac{m+1}{2x}.$$

*Proof.* Expanding the power in the left-hand side, we get

$$\left(1 + \frac{1}{x}\right)^m = \sum_{j=0}^m \binom{m}{j} \frac{1}{x^j} \geq \sum_{j=0}^1 \binom{m}{j} \frac{1}{x^j} = 1 + \frac{m}{x} \geq 1 + \frac{m+1}{2x},$$

since

$$\frac{m}{x} \geq \frac{m+1}{2x} \iff m \geq \frac{m+1}{2} \iff m \geq 1. \quad \square$$

LEMMA B.2. *For any integer  $m \geq 0$  and real  $x \geq 0$ ,*

$$(x+1)^{m+1} - x^{m+1} \leq \frac{m+1}{2} [(x+1)^m + x^m].$$

*Proof.* Expanding the powers, our inequality can be rewritten equivalently as

$$\begin{aligned} \sum_{j=0}^{m+1} \binom{m+1}{j} x^j - x^{m+1} &\leq \frac{m+1}{2} \left[ \sum_{j=0}^m \binom{m}{j} x^j + x^m \right], \\ \sum_{j=0}^{m-1} \binom{m+1}{j} x^j + (m+1)x^m &\leq \frac{m+1}{2} \left[ \sum_{j=0}^{m-1} \binom{m}{j} x^j + 2x^m \right], \\ \sum_{j=0}^{m-1} \binom{m+1}{j} x^j &\leq \frac{m+1}{2} \sum_{j=0}^{m-1} \binom{m}{j} x^j. \end{aligned}$$

Now, we can see that the above holds by bounding each term; for integers  $j = 0, 1, \dots, m-1$ ,

$$\begin{aligned} \binom{m+1}{j} &= \frac{(m+1)!}{(m+1-j)!j!} = \frac{m+1}{m+1-j} \frac{m!}{(m-j)!j!} \\ &= \frac{m+1}{m+1-j} \binom{m}{j} \leq \frac{m+1}{2} \binom{m}{j}. \end{aligned} \quad \square$$

**B.2. Proof of Lemma 4.2.** Observe that from the definition of  $A_m$  in (4.5),

$$(A_m(x))^{-1} = \frac{1}{m+1} + \frac{1}{2x},$$

which is decreasing with respect to  $m$ , and

$$\left(\frac{A_m(x)}{m+1}\right)^{-1} = 1 + \frac{m+1}{2x},$$



which is increasing with respect to  $m$ . For the remaining sequence, observe that for any integer  $m \geq 0$  and reals  $y \geq x \geq 1$ ,

$$\frac{A_m(x)}{A_m(y)} \geq \frac{A_{m+1}(x)}{A_{m+1}(y)} \iff \frac{A_{m+1}(y)}{A_m(y)} \geq \frac{A_{m+1}(x)}{A_m(x)},$$

so it is enough to show that function  $\frac{A_{m+1}(x)}{A_m(x)}$  is monotonically increasing with respect to  $x \geq 0$ . Indeed,

$$\begin{aligned} \frac{A_{m+1}(x)}{A_m(x)} &= \frac{\frac{1}{m+1} + \frac{1}{2x}}{\frac{1}{m+2} + \frac{1}{2x}} = \frac{2x(m+2) + (m+1)(m+2)}{2x(m+1) + (m+1)(m+2)} \\ &= 1 + \frac{1}{m+1} \left(1 + \frac{m+2}{2x}\right)^{-1}. \end{aligned}$$

**B.3. Proof of Lemma 4.3.** First, for (4.9), notice that it can be rewritten equivalently as

$$\frac{1}{m+1} \leq \frac{S_m(\gamma x)}{(\gamma x)^{m+1}} = \frac{1}{A_m(\gamma x)} \leq \frac{1}{A_m(\gamma)},$$

which holds, as an immediate consequence of the monotonicity of function  $A_m$  (see (4.6)), given that  $\gamma x \geq \gamma \geq 1$ . For (4.8), it is enough to prove just the special case when  $w = 1$ , i.e.,

$$(B.1) \quad \frac{\gamma^{m+1}}{A_m(\gamma)} = S_m(\gamma) \leq \frac{S_m(\gamma x + \gamma) - S_m(\gamma x)}{(x+1)^m} \leq \gamma^{m+1},$$

since then it is not difficult to check that we can recover the more general case in (4.8) by simply substituting  $\gamma := \gamma w$  and  $x := \frac{x}{w}$  into (B.1).

It is not difficult to check that (B.1) holds for  $m = 0$ , recalling that  $S_0(x) = x$  for all  $x \geq 0$ . Next, assume for the remainder of the proof that  $m \geq 1$ .

For the left-hand inequality of (B.1) first, it can be equivalently rewritten as

$$\begin{aligned} (x+1)^m \left( \frac{1}{m+1} \gamma^{m+1} + \frac{1}{2} \gamma^m \right) &\leq \frac{1}{m+1} \gamma^{m+1} [(x+1)^{m+1} - x^{m+1}] + \frac{1}{2} \gamma^m [(x+1)^m - x^m], \\ \frac{1}{2} x^m &\leq \frac{\gamma}{m+1} [(x+1)^{m+1} - (x+1)^m - x^{m+1}], \end{aligned}$$

and since  $\gamma \geq 1$ , it is sufficient to show that

$$(m+1)x^m \leq 2 [(x+1)^{m+1} - (x+1)^m - x^{m+1}]$$

and, thus, enough to show that

$$(m+1)x^m \leq 2x [(x+1)^m - x^m].$$

Now observe that the above trivially holds if  $x = 0$ , while for  $x > 0$  it can be equivalently written as

$$\frac{m+1}{2x} \leq \left(1 + \frac{1}{x}\right)^m - 1,$$

which holds due to Lemma B.1.

For the right-hand inequality of (B.1), it can be equivalently written as

$$\begin{aligned} \frac{1}{m+1} \gamma [(x+1)^{m+1} - x^{m+1}] + \frac{1}{2} [(x+1)^m - x^m] &\leq \gamma(x+1)^m, \\ 2\gamma [(x+1)^{m+1} - x^{m+1}] &\leq (m+1) [(2\gamma - 1)(x+1)^m + x^m], \\ (x+1)^{m+1} - x^{m+1} &\leq (m+1) \left[ \left(1 - \frac{1}{2\gamma}\right) (x+1)^m + \frac{1}{2\gamma} x^m \right]. \end{aligned}$$

Since  $\gamma \geq 1$ , we know that  $\frac{1}{2\gamma} \in [0, \frac{1}{2}]$ . Thus, taking into consideration that  $(x+1)^m > x^m \geq 0$ , the linear combination on the right-hand side of the above inequality is minimized for  $\frac{1}{2\gamma} = \frac{1}{2}$ . So, it is enough to show that

$$(x+1)^{m+1} - x^{m+1} \leq \frac{m+1}{2} [(x+1)^m + x^m],$$

which holds due to Lemma B.2.

**B.4. Equivalence of Theorem 4.4 and Claim 4.7.** To verify that Claim 4.7 indeed gives an equivalent restatement of Theorem 4.4, fix an arbitrary  $W \geq 1$  and observe the equivalence

$$\alpha = A_d(\gamma W) = \frac{2(d+1)\gamma W}{2\gamma W + d + 1} \iff \gamma = \frac{1}{2W} \frac{\alpha(d+1)}{d+1 - \alpha}$$

by using the definition of function  $A_d$  from (4.5). Therefore, it is not difficult to also compute that

$$\begin{aligned} \frac{d+1}{A_d(\gamma)} &= (d+1) \left( \frac{1}{d+1} + \frac{1}{2\gamma} \right) = 1 + \frac{d+1}{2} \frac{1}{\gamma} \\ &= 1 + \frac{d+1}{2} \cdot 2W \frac{d+1 - \alpha}{\alpha(d+1)} = 1 + W \left( \frac{d+1}{\alpha} - 1 \right). \end{aligned}$$

**Appendix C. Beyond polynomial latencies: Euler–Maclaurin.** Our definition of the approximate potential function in subsections 4.2 and 4.4 was based on Faulhaber’s formula (4.3) for the sum of powers of positive integers. This approach can be generalized further by considering the Euler–Maclaurin summation formula<sup>12</sup>

$$(C.1) \quad \sum_{j=0}^n f(j) = \int_0^n f(t) dt + \frac{1}{2}[f(n) + f(0)] + \sum_{j=2}^m \frac{B_j}{j!} [f^{(j-1)}(n) - f^{(j-1)}(0)] + \mathcal{R}_m$$

for any infinitely differentiable function  $f : [0, \infty) \rightarrow (0, \infty)$  (with  $f^{(j)}$  denoting the  $j$ th order derivative of  $f$ ) and integers  $n, m \geq 1$ , where  $B_j$  denotes the Bernoulli numbers we have already used in subsection 4.2 and the *error-term*  $\mathcal{R}_m$  can be bounded by

$$(C.2) \quad |\mathcal{R}_m| \leq \frac{2\zeta(m)}{(2\pi)^m} \int_0^n |f^{(m)}(t)| dt,$$

<sup>12</sup>See, e.g., [31, section 9.5] and [40].

where  $\zeta(m) = \sum_{j=1}^{\infty} \frac{1}{j^m}$  is Riemann's zeta function. Thus, if function  $f$  is such that the quantity in the right-hand side of (C.2) eventually vanishes, i.e., for any real  $x \geq 0$ ,

$$(C.3) \quad \lim_{m \rightarrow \infty} \frac{\zeta(m)}{(2\pi)^m} \int_0^x |f^{(m)}(t)| dt = 0,$$

then we can define our approximate-potential candidate function on any real  $x \geq 0$  by generalizing (C.1):

$$(C.4) \quad S(x) = S_f(x) = \int_0^x f(t) dt + \frac{1}{2}[f(x) + f(0)] + \sum_{j=2}^{\infty} \frac{B_j}{j!} [f^{(j-1)}(x) - f^{(j-1)}(0)].$$

For example, it is not difficult to see that, for any monomial  $f(x) = x^d$  of degree  $d \geq 1$ , condition (C.3) is indeed satisfied (since  $f^{(m)} = 0$  for all  $m \geq d + 1$ ), and, because also  $f^{(m)}(0) = 0$  and  $f^{(m)}(x) = \frac{d!}{(d-m)!} x^{d-m}$ , one recovers exactly (4.3) from (C.4) above.

Let us now demonstrate this general approach for latency functions  $f$  that are not polynomials. For the remainder of this section, let  $f(x) = e^x$  be an exponential delay function. Then, for any  $y \geq 0$ ,

$$\lim_{m \rightarrow \infty} \frac{\zeta(m)}{(2\pi)^m} \int_0^y |f^{(m)}(t)| dt = (e^y - 1) \lim_{m \rightarrow \infty} \frac{\zeta(m)}{(2\pi)^m} = 0,$$

since  $\lim_{m \rightarrow \infty} \zeta(m) = 1$  and  $\lim_{m \rightarrow \infty} (2\pi)^m = \infty$ . Thus, condition (C.3) is satisfied, and we can define from (C.4)

$$\begin{aligned} S(x) &= \int_0^x e^t dt + \frac{1}{2}[e^x + e^0] + \sum_{j=2}^{\infty} \frac{B_j}{j!} [e^x - e^0] \\ &= (e^x - 1) - \frac{1}{2}(e^x - 1) + \sum_{j=2}^{\infty} \frac{B_j}{j!} (e^x - 1) + e^x \\ &= (e^x - 1) \sum_{j=0}^{\infty} \frac{B_j}{j!} + e^x. \end{aligned}$$

But since for the integer value  $x = 1$  we know that  $S(1) = \sum_{j=0}^1 f(j) = 1 + e$ , it must be that

$$e + 1 = (e^x - 1) \sum_{j=0}^{\infty} \frac{B_j}{j!} + e \iff \sum_{j=0}^{\infty} \frac{B_j}{j!} = \frac{1}{e - 1}.$$

So, we finally have that

$$S(x) = (e^x - 1) \frac{1}{e - 1} + e^x = \frac{e^{x+1} - 1}{e - 1}.$$

From this, for all reals  $x \geq 0$ ,  $w > 0$  we compute

$$(C.5) \quad \frac{S(x+w) - S(x)}{wf(x+w)} = \frac{1}{e-1} \frac{e^{x+w+1} - e^{x+1}}{we^{x+w}} = \frac{e}{e-1} \frac{1 - e^{-w}}{w},$$

which does not depend on  $x$ . Thus, from (4.1) in Lemma 4.1 we deduce that *exact* pure Nash equilibria always exist for weighted congestion games with exponential

latencies. The function  $S$  we defined in (C.5) essentially serves as a weighted potential [43]; its global minimum is a pure Nash equilibrium. Notice here that these results regarding exponential latency functions were already known because of the work of Panagopoulou and Spirakis [45].

**Appendix D. Social optimum is a  $(d + 1)$ -approximate equilibrium.** In this section we show that the socially optimum solution is itself a  $(d + 1)$ -approximate equilibrium, where  $d$  is the maximum degree of the polynomial latency functions. We must mention here that the approximation factor on its own, i.e.,  $d + 1$ , does not constitute a novel contribution: existence of  $(d + 1)$ -approximate equilibria was already known because of the work of Harks and Klimm [33]. However, the new element in Theorem D.1 below is that this can be achieved by all optimum solutions.

Related to this, we would like to emphasize that if we only cared about showing the existence of *an* optimal solution that is a  $(d + 1)$ -approximate equilibrium, this would have been an immediate corollary of our main upper bound result: by simply setting  $\alpha = d + 1$  in Theorem 4.4 we get exactly what we want. However, the following theorem demonstrates the stronger statement that *all* social cost minimizers have the property we want.

**THEOREM D.1.** *Consider any weighted congestion game with polynomial latency functions of maximum degree  $d$ , and let  $\mathbf{s}^*$  be a strategy profile that minimizes social cost. Then  $\mathbf{s}^*$  is a  $(d + 1)$ -approximate pure Nash equilibrium. As an immediate consequence, the PoS of  $(d + 1)$ -approximate Nash equilibria is 1.*

*Proof.* Let  $c$  be an arbitrary cost function of maximum degree  $d$  with nonnegative coefficients, i.e.,  $c(x) = \sum_{j=0}^d a_j x^j$ , with  $a_j \geq 0$  for all  $j$ . We will first show that for all  $w > 0$  and  $x \geq 0$ ,

$$(D.1) \quad w \cdot c(x + w) \leq (x + w) \cdot c(x + w) - x \cdot c(x) \leq (d + 1) \cdot w \cdot c(x + w).$$

To this end, with  $z = \frac{x}{w}$ , we get

$$\begin{aligned} (x + w) \cdot c(x + w) - x \cdot c(x) &= \sum_{j=0}^d a_j \cdot [(x + w)^{j+1} - x^{j+1}] \\ &= \sum_{j=0}^d a_j \cdot w^{j+1} [(1 + z)^{j+1} - z^{j+1}] \\ &= \sum_{j=0}^d a_j \cdot w^{j+1} \left[ \sum_{k=0}^j \binom{j+1}{k} z^k \right] \end{aligned}$$

and

$$\begin{aligned} w \cdot c(x + w) &= \sum_{j=0}^d a_j \cdot w(x + w)^j \\ &= \sum_{j=0}^d a_j \cdot w^{j+1} \left[ \sum_{k=0}^j \binom{j}{k} z^k \right]. \end{aligned}$$

Clearly,  $\binom{j+1}{k} \geq \binom{j}{k}$  for all integers  $j \in [0, d], k \in [0, j]$ , which immediately implies

the first inequality in (D.1). To see the second inequality, observe that

$$\frac{\binom{j+1}{k}}{\binom{j}{k}} = \frac{j+1}{j+1-k} \leq j+1 \leq d+1.$$

Since  $\mathbf{s}^*$  minimizes social cost, for all players  $i \in [n]$  and strategies  $s_i \in S_i$ ,

$$C(\mathbf{s}^*) \leq C(s_i, \mathbf{s}_{-i}^*).$$

Denoting  $y_e = \sum_{j \in [n] \setminus \{i\}: e \in s_j^*} w_j$ , from (D.1), we get

$$\begin{aligned} 0 &\leq C(s_i, \mathbf{s}_{-i}^*) - C(\mathbf{s}^*) \\ &= \sum_{e \in s_i \setminus s_i^*} [(y_e + w_i)c_e(y_e + w_i) - y_e c_e(y_e)] \\ &\quad - \sum_{e \in s_i^* \setminus s_i} [(y_e + w_i)c_e(y_e + w_i) - y_e c_e(y_e)] \\ &= \sum_{e \in s_i} [(y_e + w_i)c_e(y_e + w_i) - y_e c_e(y_e)] - \sum_{e \in s_i^*} [(y_e + w_i)c_e(y_e + w_i) - y_e c_e(y_e)] \\ &\leq (d+1) \sum_{e \in s_i} w_i \cdot c_e(y_e + w_i) - \sum_{e \in s_i^*} w_i \cdot c_e(y_e + w_i) \\ &= (d+1)C_i(s_i, \mathbf{s}_{-i}^*) - C_i(\mathbf{s}^*), \end{aligned}$$

or, equivalently,  $C_i(\mathbf{s}^*) \leq (d+1)C_i(s_i, \mathbf{s}_{-i}^*)$ . So  $\mathbf{s}^*$  is a  $(d+1)$ -approximate Nash equilibrium.  $\square$

**Acknowledgment.** We thank the anonymous reviewers for their careful and thorough reading of our manuscript and for their valuable feedback.

#### REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 10th ed., Dover, New York, 1964, <http://people.math.sfu.ca/~cbm/aands/>.
- [2] H. ACKERMANN, H. RÖGLIN, AND B. VÖCKING, *On the impact of combinatorial structure on congestion games*, J. ACM, 55 (2008), pp. 1–22, <https://doi.org/10.1145/1455248.1455249>.
- [3] S. ALAND, D. DUMRAUF, M. GAIRING, B. MONIEN, AND F. SCHOPPMANN, *Exact price of anarchy for polynomial congestion games*, SIAM J. Comput., 40 (2011), pp. 1211–1233, <https://doi.org/10.1137/090748986>.
- [4] S. ALBERS, *On the value of coordination in network design*, SIAM J. Comput., 38 (2009), pp. 2273–2302, <https://doi.org/10.1137/070701376>.
- [5] E. ANSHELEVICH, A. DASGUPTA, J. KLEINBERG, É. TARDOS, T. WEXLER, AND T. ROUGHGARDEN, *The price of stability for network design with fair cost allocation*, SIAM J. Comput., 38 (2008), pp. 1602–1623, <https://doi.org/10.1137/070680096>.
- [6] B. AWERBUCH, Y. AZAR, AND A. EPSTEIN, *The price of routing unsplittable flow*, SIAM J. Comput., 42 (2013), pp. 160–177, <https://doi.org/10.1137/070702370>.
- [7] J. BERNOULLI, *Ars conjectandi, opus posthumum. Accedit Tractatus de seriebus infinitis, et epistola gallicé scripta de ludo pilae reticularis*, impensis Thurnisiorum, fratrum, Basileæ, 1713, <https://books.google.de/books?id=ebdcNfPz8l8C>.
- [8] K. BHAWALKAR, M. GAIRING, AND T. ROUGHGARDEN, *Weighted congestion games: The price of anarchy, universal worst-case examples, and tightness*, ACM Trans. Econ. Comput., 2 (2014), 14, <https://doi.org/10.1145/2629666>.
- [9] V. BILÒ, *A unifying tool for bounding the quality of non-cooperative solutions in weighted congestion games*, Theory Comput. Syst., 62 (2018), pp. 1288–1317, <https://doi.org/10.1007/s00224-017-9826-1>.

- [10] V. BILÒ, I. CARAGIANNIS, A. FANELLI, AND G. MONACO, *Improved lower bounds on the price of stability of undirected network design games*, *Theory Comput. Syst.*, 52 (2013), pp. 668–686, <https://doi.org/10.1007/s00224-012-9411-6>.
- [11] V. BILÒ AND C. VINCI, *On the impact of singleton strategies in congestion games*, in *Proceedings of the 25th Annual European Symposium on Algorithms*, Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, Wadern, Germany, 2017, 17, <https://doi.org/10.4230/lipics.esa.2017.17>.
- [12] V. BILÒ, M. FLAMMINI, AND L. MOSCARDELLI, *The price of stability for undirected broadcast network design with fair cost allocation is constant*, *Games Econom. Behavior*, to appear, <https://doi.org/10.1016/j.geb.2014.09.010>.
- [13] I. CARAGIANNIS, M. FLAMMINI, C. KAKLAMANIS, P. KANELLOPOULOS, AND L. MOSCARDELLI, *Tight bounds for selfish and greedy load balancing*, *Algorithmica*, 61 (2011), pp. 606–637, <https://doi.org/10.1007/s00453-010-9427-8>.
- [14] I. CARAGIANNIS, A. FANELLI, N. GRAVIN, AND A. SKOPALIK, *Efficient computation of approximate pure Nash equilibria in congestion games*, in *Proceedings of the 52nd IEEE Annual Symposium on Foundations of Computer Science*, Palm Springs, CA, 2011, <https://doi.org/10.1109/focs.2011.50>.
- [15] I. CARAGIANNIS, A. FANELLI, N. GRAVIN, AND A. SKOPALIK, *Approximate pure Nash equilibria in weighted congestion games: Existence, efficient computation, and structure*, *ACM Trans. Econ. Comput.*, 3 (2015), 2, <https://doi.org/10.1145/2614687>.
- [16] H.-L. CHEN AND T. ROUGHGARDEN, *Network design with weighted players*, *Theory Comput. Syst.*, 45 (2008), 302, <https://doi.org/10.1007/s00224-008-9128-8>.
- [17] G. CHRISTODOULOU AND M. GAIRING, *Price of stability in polynomial congestion games*, *ACM Trans. Econ. Comput.*, 4 (2016), 10, <https://doi.org/10.1145/2841229>.
- [18] G. CHRISTODOULOU AND E. KOUTSOPIAS, *The price of anarchy of finite congestion games*, in *Proceedings of the 37th Annual ACM Symposium on Theory of Computing (STOC '05)*, ACM, New York, 2005, pp. 67–73, <https://doi.org/10.1145/1060590.1060600>.
- [19] G. CHRISTODOULOU AND E. KOUTSOPIAS, *On the price of anarchy and stability of correlated equilibria of linear congestion games*, in *Algorithms—ESA 2005*, Springer, Berlin, 2005, pp. 59–70.
- [20] G. CHRISTODOULOU, E. KOUTSOPIAS, AND P. G. SPIRAKIS, *On the performance of approximate equilibria in congestion games*, *Algorithmica*, 61 (2011), pp. 116–140, <https://doi.org/10.1007/s00453-010-9449-2>.
- [21] G. CHRISTODOULOU, M. GAIRING, Y. GIANNAKOPOULOS, AND P. G. SPIRAKIS, *The price of stability of weighted congestion games*, in *45th International Colloquium on Automata, Languages, and Programming, ICALP 2018*, Prague, Czech Republic, 2018, 150, <https://doi.org/10.4230/LIPIcs.ICALP.2018.150>.
- [22] J. H. CONWAY AND R. K. GUY, *The Book of Numbers*, Springer, New York, 1996, <https://doi.org/10.1007/978-1-4612-4072-3>.
- [23] J. DUNKEL AND A. S. SCHULZ, *On the complexity of pure-strategy Nash equilibria in congestion and local-effect games*, *Math. Oper. Res.*, 33 (2008), pp. 851–868.
- [24] A. FABRIKANT, C. PAPADIMITRIOU, AND K. TALWAR, *The complexity of pure Nash equilibria*, in *Proceedings of the 36th Annual ACM Symposium on Theory of Computing, STOC '04*, ACM, New York, 2004, pp. 604–612, <https://doi.org/10.1145/1007352.1007445>.
- [25] J. FAULHABER, *Academia Algebrae: Darinnen die miraculosische Inventiones zu den höchsten Cossen weiters continuirt und profitiert werden*, Johann Ulrich Schönigs, Augspurg, 1631, [https://books.google.de/books?id=0pw\\_AAAAcAAJ](https://books.google.de/books?id=0pw_AAAAcAAJ).
- [26] M. FELDOTTO, M. GAIRING, G. KOTSIALOU, AND A. SKOPALIK, *Computing approximate pure Nash equilibria in Shapley value weighted congestion games*, in *Web and Internet Economics 2017*, Springer, Cham, 2017, pp. 191–204, [https://doi.org/10.1007/978-3-319-71924-5\\_14](https://doi.org/10.1007/978-3-319-71924-5_14).
- [27] A. FIAT, H. KAPLAN, M. LEVY, S. OLONETSKY, AND R. SHABO, *On the price of stability for designing undirected networks with fair cost allocations*, in the *33rd International Colloquium on Automata, Languages and Programming (ICALP)*, M. Bugliesi, B. Preneel, V. Sassone, and I. Wegener, eds., Springer, Berlin, 2006, pp. 608–618, [https://doi.org/10.1007/11786986\\_53](https://doi.org/10.1007/11786986_53).
- [28] D. FOTAKIS, S. KONTOGIANNIS, AND P. SPIRAKIS, *Selfish unsplittable flows*, *Theoret. Comput. Sci.*, 348 (2005), pp. 226–239.
- [29] D. FOTAKIS, S. KONTOGIANNIS, E. KOUTSOPIAS, M. MAVRONICOLAS, AND P. SPIRAKIS, *The structure and complexity of Nash equilibria for a selfish routing game*, *Theoret. Comput. Sci.*, 410 (2009), pp. 3305–3326, <https://doi.org/10.1016/j.tcs.2008.01.004>.

- [30] M. GOEMANS, V. MIRROKNI, AND A. VETTA, *Sink equilibria and convergence*, in Proceedings of the 46th Annual IEEE Symposium on Foundations of Computer Science (FOCS'05), Pittsburgh, PA, 2005, pp. 142–151, <https://doi.org/10.1109/SFCS.2005.68>.
- [31] R. L. GRAHAM, D. E. KNUTH, AND O. PATASHNIK, *Concrete Mathematics: A Foundation for Computer Science*, Addison-Wesley, Reading, MA, 1989.
- [32] C. HANSKNECHT, M. KLIMM, AND A. SKOPALIK, *Approximate pure Nash equilibria in weighted congestion games*, in Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2014), Leibniz International Proceedings in Informatics (LIPIcs) 28, K. Jansen, J. D. P. Rolim, N. R. Devanur, and C. Moore, eds., Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, Wadern, Germany, 2014, pp. 242–257, <https://doi.org/10.4230/LIPIcs.APPROX-RANDOM.2014.242>.
- [33] T. HARKS AND M. KLIMM, *On the existence of pure Nash equilibria in weighted congestion games*, *Math. Oper. Res.*, 37 (2012), pp. 419–436, <https://doi.org/10.1287/moor.1120.0543>.
- [34] T. HARKS, M. KLIMM, AND R. H. MÖHRING, *Characterizing the existence of potential functions in weighted congestion games*, *Theory Comput. Syst.*, 49 (2011), pp. 46–70, <https://doi.org/10.1007/s00224-011-9315-x>.
- [35] T. HARKS, M. KLIMM, AND R. H. MÖHRING, *Strong equilibria in games with the lexicographical improvement property*, *Internat. J. Game Theory*, 42 (2012), pp. 461–482, <https://doi.org/10.1007/s00182-012-0322-1>.
- [36] C. G. J. JACOBI, *De usu legitimo formulae summatoriae Maclauriniana*, *J. Reine Angew. Math.*, 12 (1834), pp. 263–272, <http://www.digizeitschriften.de/dms/img/?PID=GDZPPN00214008X>.
- [37] D. E. KNUTH, *Johann Faulhaber and sums of powers*, *Math. Comp.*, 61 (1993), pp. 277–277, <https://doi.org/10.1090/s0025-5718-1993-1197512-7>.
- [38] E. KOUTSOPIAS AND C. PAPADIMITRIOU, *Worst-case equilibria*, in Proceedings of the 16th Annual ACM Symposium on Theoretical Aspects of Computer Science (STACS '99), ACM, New York, 1999, pp. 404–413.
- [39] E. LEE AND K. LIGETT, *Improved bounds on the price of stability in network cost sharing games*, in Proceedings of the 14th ACM Conference on Electronic Commerce (EC '13), ACM, New York, 2013, pp. 607–620, <https://doi.org/10.1145/2492002.2482562>.
- [40] D. H. LEHMER, *On the maxima and minima of Bernoulli polynomials*, *Amer. Math. Monthly*, 47 (1940), pp. 533–538, <https://doi.org/10.2307/2303833>.
- [41] L. LIBMAN AND A. ORDA, *Atomic resource sharing in noncooperative networks*, *Telecommunication Syst.*, 17 (2001), pp. 385–409, <https://doi.org/10.1023/A:1016770831869>.
- [42] D. S. MITRINOVIĆ, *Analytic Inequalities*, Springer, Berlin, Heidelberg, 1970, <https://doi.org/10.1007/978-3-642-99970-3>.
- [43] D. MONDERER AND L. S. SHAPLEY, *Potential games*, *Games Econom. Behavior*, 14 (1996), pp. 124–143.
- [44] N. NISAN, T. ROUGHGARDEN, É. TARDOS, AND V. VAZIRANI, EDS., *Algorithmic Game Theory*, Cambridge University Press, Cambridge, UK, 2007.
- [45] P. N. PANAGOPOULOU AND P. G. SPIRAKIS, *Algorithms for pure Nash equilibria in weighted congestion games*, *J. Experimental Algorithmics*, 11 (2007), 2.7, <https://doi.org/10.1145/1187436.1216584>.
- [46] R. W. ROSENTHAL, *A class of games possessing pure-strategy Nash equilibria*, *Internat. J. Game Theory*, 2 (1973), pp. 65–67.
- [47] R. W. ROSENTHAL, *The network equilibrium problem in integers*, *Networks*, 3 (1973), pp. 53–59.
- [48] T. ROUGHGARDEN, *Intrinsic robustness of the price of anarchy*, *J. ACM*, 62 (2015), 32, <https://doi.org/10.1145/2806883>.
- [49] T. ROUGHGARDEN AND É. TARDOS, *How bad is selfish routing?*, *J. ACM*, 49 (2002), pp. 236–259, <https://doi.org/10.1145/506147.506153>.
- [50] T. ROUGHGARDEN AND É. TARDOS, *Bounding the inefficiency of equilibria in nonatomic congestion games*, *Games Econom. Behavior*, 47 (2004), pp. 389–403, <https://doi.org/10.1016/j.geb.2003.06.004>.
- [51] A. S. SCHULZ AND N. S. MOSES, *On the performance of user equilibria in traffic networks*, in Proceedings of the 14th Annual ACM–SIAM Symposium on Discrete Algorithms (Baltimore, MD), ACM, New York, SIAM, Philadelphia, 2003, pp. 86–87.